

2004

PREPRINT 277

Michel Janssen and Matthew Mecklenburg

**Electromagnetic Models of the Electron and
the Transition from Classical to Relativistic
Mechanics**

Electromagnetic Models of the Electron and the Transition from Classical to Relativistic Mechanics*

Michel Janssen and Matthew Mecklenburg

1. Introduction

“Special relativity killed the classical dream of using the energy-momentum-velocity relations as a means of probing the dynamical origins of [the mass of the electron]. The relations are purely kinematical” (Pais 1982, 159). This perceptive comment comes from a section on the pre-relativistic notion of electromagnetic mass in ‘*Subtle is the Lord ...*’, Abraham Pais’ highly acclaimed biography of Albert Einstein. By ‘kinematical’ Pais meant something like ‘completely independent of the details of the dynamics’. In this paper we examine the classical dream referred to by Pais from the vantage point of relativistic continuum mechanics.

There were actually two such dreams in the years surrounding the advent of special relativity. Like Einstein’s theory, both dreams originated in the electrodynamics of moving bodies developed in the 1890s by the Dutch physicist Hendrik Antoon Lorentz. Both took the form of concrete models of the electron. Even these models were similar. Yet they were part of fundamentally different programs competing with one another in the years around 1905. One model, due to the German theoretician Max Abraham (1902a), was part of a revolutionary effort to substitute the laws of electrodynamics for those of Newtonian mechanics as the fundamental laws of physics. The other model, adapted from Abraham’s by Lorentz (1904b) and fixed up by the French mathematician Henri Poincaré (1906), was part of the attempt to provide a general explanation for the absence of any signs of ether drift, the elusive 19th-century medium thought to carry light waves and electromagnetic fields. A choice had to be made between the objectives of Lorentz and Abraham. One could not eliminate all signs of the earth’s motion through the ether and reduce all physics to electrodynamics at the same time. Special relativity was initially conflated with Lorentz’s theory because it too seemed to focus on the undetectability of motion at the expense of electromagnetic purity. The theories of

* To appear in a volume edited by Jesper Lützen based on the proceedings of *The Interaction between Mathematics, Physics and Philosophy from 1850 to 1940*, a conference held at the Carlsberg Academy, Copenhagen, September 26–28, 2002.

Lorentz and Einstein agreed in all their empirical predictions, including those for the velocity-dependence of electron mass, even though special relativity was not wedded to any particular model of the electron. For a while there was a third electron model, a variant on Lorentz's proposed independently by Alfred Bucherer (1904, 57–60; 1905) and Paul Langevin (1905). At the time, the acknowledged arbiter between these models and the broader theories (perceived to be) attached to them was a series of experiments by Walter Kaufmann and others on the deflection of high-speed electrons in β -radiation and cathode rays by electric and magnetic fields for the purpose of determining the velocity-dependence of their mass.¹

As appropriate for reveries, neither Lorentz's nor Abraham's dream about the nature and structure of the electron lasted long. They started to fade a few years after Einstein's formulation of special relativity, even though the visions that inspired them lingered on for quite a while. Lorentz went to his grave clinging to the notion of an ether hidden from view by the Lorentz-invariant laws governing the phenomena. Abraham's electromagnetic vision was pursued well into the 1920s by kindred spirits such as Gustav Mie (1912a, b; 1913). By then mainstream physics had long moved on. The two dreams, however, did not evaporate without a trace. They played a decisive role in the development of relativistic mechanics.² It is no coincidence therefore that relativistic continuum mechanics will be central to our analysis in this paper. The development of the new mechanics effectively began with the non-Newtonian transformation laws for force and mass introduced by Lorentz in the 1890s. It continued with the introduction of electromagnetic momentum and electromagnetic mass by Abraham (1902a, b; 1903; 1904; 1905, 1909) in the wake of the proclamation of the electromagnetic view of nature by Willy Wien (1900). Einstein (1907b), Max Planck (1906a, 1908), Hermann Minkowski (1908), Arnold Sommerfeld (1910a, b), and Gustav Herglotz (1910, 1911)—the last three champions of the electromagnetic program³—all contributed to its

¹ See (Kaufmann 1901a, b, c; 1902a, b; 1903; 1905; 1906a, b; 1907). For references to later experiments, see, e.g., (Pauli 1921, 83). Our paper will not touch on the intricacies of the actual experiments. Those are covered in (Cushing 1981). See also (Hon 1995).

² Moreover, the concept of "Poincaré pressure" introduced to stabilize Lorentz's electron (see below) resurfaced in a theory of Einstein (1919), which is enjoying renewed interest (Earman 2003), as well as in other places (see, e.g., Grøn 1985, 1988).

³ See (McCormach 1970, 490) for a discussion of the development of Sommerfeld's attitude toward the electromagnetic program and special relativity. On Minkowski and the electromagnetic program, see (Galison 1979), (Pyenson 1985, Ch. 4), and, especially, (Corry 1997).

further development in a proper relativistic setting. These efforts culminated in a seminal paper by Max Laue (1911a) and were enshrined in the first textbook on relativity published later that year (Laue 1911b).

There already exists a voluminous literature on the various aspects of this story. We shall freely draw and build on that literature. One of us has written extensively on the development of Lorentz's research program in the electrodynamics of moving bodies (Janssen 1995, 2002b; Janssen and Stachel 2004).⁴ The canonical source for the electromagnetic view of nature is still (McCormmach 1970), despite its focus on Lorentz whose attitude toward the electromagnetic program was ambivalent (cf. Lorentz 1900; 1905, 93–101; 1915, secs. 178–186). His work formed its starting point and he was sympathetic to the program, but never a strong advocate of it. (Goldberg 1970) puts the spotlight on the program's undisputed leader, Max Abraham. (Pauli 1921, Ch. 5) is a good source for the degenerative phase of the electromagnetic program in the 1910s.⁵ For a concise overview of the rise and fall of the electromagnetic program, see Ch. 8 of (Kragh 1999), aptly titled "A Revolution that Failed."

Another important source for the electromagnetic program is Ch. 5 in (Pyenson 1985), which discusses a seminar on electron theory held in Göttingen in the summer semester of 1905. Minkowski was one of four instructors of this course. The other three were David Hilbert, Emil Wiechert, and Herglotz, who had already published on electron theory (Herglotz 1903). Max Laue audited the seminar as a postdoc. The syllabus for the seminar lists papers by Lorentz (1904a, b), Abraham (1903), Karl Schwarzschild (1903a, b, c), and Sommerfeld (1904a, b; 1905). This seminar gives a good indication of how active and cutting edge this research area was at the time. Further evidence of this vitality is provided by debates in the literature of the day such as those between Bucherer (1907; 1908a, b) and Ebenezer Cunningham (1907, 1908)⁶ and between Einstein (1907a) and Paul Ehrenfest (1906, 1907) over various points concerning these electron models. The roll call of researchers active in this area also included the Italian mathematician Tullio Levi-Civita (1907, 1909).⁷ One may even get the impression that in the early 1900s the

⁴ See also (Darrigol 2000). We refer to (Janssen 1995, 2002b) for references to and discussion of earlier literature on this topic.

⁵ For more recent commentary, see (Corry 1999).

⁶ For brief discussions, see (Balazs 1972, 29–30) and (Warwick 2003, Ch. 8, especially 413–414). We also refer to Warwick's work for British reactions to the predominantly German developments discussed in our paper. See, e.g., (Warwick 2003, 384) for comments by James Jeans on electromagnetic mass.

⁷ For brief discussion, see (Balazs 1972, 30)

journals were flooded with papers on electron models. We wonder, for instance, whether the book on electron theory by Bucherer (1904) was not originally written as a long journal article, which was rejected, given its similarity to earlier articles by Abraham, Lorentz, Schwarzschild, and Sommerfeld.

The saga of the Abraham, Lorentz, and Bucherer-Langevin electron models and their changing fortunes in the laboratories of Kaufmann, Bucherer, and others has been told admirably by Arthur I. Miller (1981, secs. 1.8–1.14, 7.4, and 12.4). Miller (1973) is also responsible for a detailed analysis of the classic paper by Poincaré (1906) that introduced what came to be known as “Poincaré pressure” to stabilize Lorentz’s purely electromagnetic electron.⁸ Fritz Rohrlich (1960, 1965, 1973) has given a particularly insightful analysis of the electron model of Lorentz and Poincaré.⁹ The model is also covered concisely and elegantly in volume two of the Feynman lectures (Feynman 1964, Ch. 28).

Given how extensively this episode has been discussed in the literature, the number of sources covering its denouement with the formulation of Laue’s relativistic continuum mechanics is surprisingly low. Max Jammer does not discuss relativistic continuum mechanics at all in his classic monograph on the development of the concept of mass (cf. Jammer 1997, Chs. 11–13). And although Miller prominently discusses Laue’s work, both in (Miller 1973, sec. 7.5) and in the concluding section of his book (Miller 1981, sec. 12.5.8), he does not give it the central place that in our opinion it deserves. To bring out the importance of Laue’s work, we show right from the start how the kind of spatially extended systems studied by Abraham, Lorentz, and Poincaré can be dealt with in special relativity. We shall use modern notation and modern units throughout and give self-contained derivations of all results. Our treatment of these electron models follows the analysis of the experiments of Trouton and Noble in (Janssen 1995, 2002b, 2003), which was inspired in part by the discussion in (Norton 1992) of the importance of Laue’s relativistic mechanics for the development of Gunnar Nordström’s special-relativistic theory of gravity. The focus on the conceptual changes in mechanics that accompanied the transition from classical to relativistic kinematics was inspired in part by the work of Jürgen Renn and his collaborators on pre-classical mechanics (Damerow et al. 2004).

⁸ We have benefited from (annotated) translations of Poincaré’s paper by Schwartz (1971, 1972) and Kilmister (1970), as well as from the translation of passages from (Poincaré 1905), the short version of (Poincaré 1906), in (Keswani and Kilmister 1983).

⁹ See also (Yaghjian 1992). (Pais 1972) is another useful source.

2. Energy-momentum-mass-velocity relations.

2.1. Special relativity. In special relativity, the relations between energy, momentum, mass, and velocity of a system are encoded in the transformation properties of its four-momentum. This quantity combines the energy U and the three components of the ordinary momentum \mathbf{P} :¹⁰

$$P^\mu = \left(\frac{U}{c}, \mathbf{P} \right) \quad (2.1)$$

(where c is the velocity of light). In the system's rest frame, with coordinates $x_0^\mu = (ct_0, x_0, y_0, z_0)$, the four-momentum reduces to:

$$P_0^\mu = \left(\frac{U_0}{c}, 0, 0, 0 \right), \quad (2.2)$$

i.e., $\mathbf{P}_0 = 0$. The system's rest mass is defined as $m_0 \equiv U_0/c^2$.

We transform P_0^μ from the x_0^μ -frame to some new x^μ -frame, assuming that P_0^μ transforms as a four-vector under Lorentz transformations. Let the two frames be related by the Lorentz transformation $x^\mu = \Lambda^\mu{}_\nu x_0^\nu$.¹¹ Since, in general, the four-momentum does *not* transform as a four-vector, the Lorentz transform of P_0^μ will, in general, not be the four-momentum in the x^μ -frame. We therefore cautiously write the result of the transformation with an asterisk:

$$P^{*\mu} = \Lambda^\mu{}_\nu P_0^\nu. \quad (2.3)$$

Without loss of generality we can focus on the special case in which the motion of the x^μ -frame with respect to the x_0^μ -frame is with velocity v in the x -direction. The matrix for this transformation is:

$$\Lambda^\mu{}_\nu = \begin{pmatrix} \gamma & \gamma\beta & 0 & 0 \\ \gamma\beta & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (2.4)$$

¹⁰ The letter U rather than E is used for energy to avoid confusion with the electric field. We shall be using SI units throughout. For conversion to other units, see, e.g., (Jackson 1975, 817–819).

¹¹ The transformation matrices $\Lambda^\mu{}_\nu$ satisfy $\Lambda^\mu{}_\rho \Lambda^\nu{}_\sigma \eta^{\rho\sigma} = \eta^{\mu\nu}$, the defining equation for Lorentz transformations, where $\eta^{\mu\nu} \equiv \text{diag}(1, -1, -1, -1)$ is the standard diagonal Minkowski metric.

with $\gamma \equiv 1/\sqrt{1-\beta^2}$ and $\beta \equiv v/c$. In that case,

$$P^{*\mu} = \left(\gamma \frac{U_0}{c}, \gamma \beta \frac{U_0}{c}, 0, 0 \right) = (\gamma m_0 c, \gamma m_0 \mathbf{v}). \quad (2.5)$$

If the four-momentum of the system *does* transform as a four-vector, $P^{*\mu}$ in eq. (2.5) is equal to P^μ in eq. (2.1) and we can read off the following relations between energy, momentum, mass, and velocity from these two equations:

$$U = \gamma U_0 = \gamma m_0 c^2, \quad \mathbf{P} = \gamma m_0 \mathbf{v}. \quad (2.6)$$

Eqs. (2.6) hold for a relativistic point particle with rest mass m_0 . Its four-momentum is given by

$$P^\mu = m_0 u^\mu = m_0 \frac{dx^\mu}{d\tau} = \gamma m_0 \frac{dx^\mu}{dt}. \quad (2.7)$$

Since $u^\mu \equiv dx^\mu/d\tau$ is the four-velocity, this is clearly a four-vector. The relation

between proper time τ , arc length s , and coordinate time t is given by

$d\tau = ds/c = dt/\gamma$.¹² If the particle is moving with velocity \mathbf{v} , $dx^\mu/dt = (c, \mathbf{v})$ and eq. (2.7) becomes:

$$P^\mu = (\gamma m_0 c, \gamma m_0 \mathbf{v}). \quad (2.8)$$

Eqs. (2.6) also hold for spatially extended *closed* systems, i.e., systems described by an energy-momentum tensor $T^{\mu\nu}$ with a vanishing four-divergence, i.e., $\partial_\nu T^{\mu\nu} = 0$.¹³ The energy-momentum tensor brings together the following quantities. The component T^{00} gives the energy density; T^{i0}/c the components of the momentum density; cT^{0i} the components of the energy flow density;¹⁴ and T^{ij} the components of the momentum flow density, or, equivalently, the stresses.¹⁵ The standard definition of the four-momentum of

¹² From $ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu = (c^2 - v^2) dt^2$ it follows that $ds = c\sqrt{1-v^2/c^2} dt = c dt/\gamma$.

¹³ Here and in the rest of the paper summation over repeated indices is implied; ∂_μ stands for $\partial/\partial x^\mu$.

¹⁴ The energy-momentum tensor is typically symmetric. In that case, $T^{i0} = T^{0i}$, which means that the momentum density (T^{i0}/c) equals the energy flow density (cT^{0i}) divided by c^2 . As was first noted by Planck, this is one way of expressing the inertia of energy, $E = mc^2$.

¹⁵ Which is why $T^{\mu\nu}$ is also known as the stress-energy tensor or the stress-energy-momentum tensor

a spatially extended (not necessarily closed) system described by the (not necessarily divergence-free) energy-momentum tensor $T^{\mu\nu}$ is:

$$P^\mu \equiv \frac{1}{c} \int T^{\mu 0} d^3x. \quad (2.9)$$

Before the advent of relativity, this equation was written as a pair of separate equations:

$$U = \int u d^3x, \quad \mathbf{P} = \int \mathbf{p} d^3x, \quad (2.10)$$

where u and \mathbf{p} are the energy density and the momentum density, respectively. Definition (2.9) is clearly not manifestly Lorentz invariant. The space integrals of $T^{\mu 0}$ in the x^μ -frame are integrals in space-time over a three-dimensional hyperplane of simultaneity in that frame. A Lorentz transformation does not change the hyperplane over which the integration is to be carried out. A hyperplane of simultaneity in the x^μ -frame is *not* a hyperplane of simultaneity in any frame moving with respect to it. From these last three observations, it follows that the Lorentz transforms of the space integrals in eq. (2.9) will not be space integrals in the new frame. But then how can these Lorentz transforms ever be the four-momentum in the new frame? The answer to this question is that *if the system is closed* (i.e., if $\partial_\nu T^{\mu\nu} = 0$), it does not matter over which hyperplane the integration is done. The integrals of the relevant components of $T^{\mu\nu}$ over any hyperplane extending to infinity will all give the same values. So for closed systems a Lorentz transformation does map the four-momentum in one frame to a quantity that is equal to the four-momentum in the new frame even though these two quantities are defined as integrals over different hyperplanes.¹⁶

The standard definition of four-momentum can be replaced by a manifestly Lorentz-invariant one. First note that the space integrals of $T^{\mu 0}$ in the x^μ -frame can be written in a manifestly covariant form as¹⁷

$$P^\mu = \frac{1}{c} \int \delta(\eta_{\rho\sigma} x^\rho n^\sigma) T^{\mu\nu} n_\nu d^4x, \quad (2.11)$$

¹⁶ See (Rohrlich 1965, 89–90, 279–281) or (Janssen 1995, sec. 2.1.3) for the details of the proof, which is basically an application of the obvious generalization of Gauss' theorem (which says that for any vector field \mathbf{A} , $\oint \mathbf{A} \cdot d\mathbf{S} = \int \text{div} \mathbf{A} d^3x$) from three to four dimensions.

¹⁷ This way of writing P^μ was suggested to me by Serge Rudaz. See (Janssen 2002b, 440–441, note; 2003, 47) for a more geometrical way of stating the argument below.

where n^μ is a unit vector in the time direction in the x^μ -frame. In that frame n^μ has components $(1, 0, 0, 0)$. The delta function picks out hyperplanes of simultaneity in the x^μ -frame. The standard definition (2.9) of four-momentum can, of course, be written in the form of eq. (2.11) in any frame, but that requires a different choice of n^μ in each one. This is just a different way of saying what we said before: under the standard definition (2.9), the result of transforming P^μ in the x^μ -frame to some new frame will *not* be the four-momentum in the new frame unless the system happens to be closed. If, however, we take the unit vector n^μ in eq. (2.11) to be some *fixed* timelike vector—typically the unit vector in the time direction in the system’s rest frame¹⁸— and take eq. (2.11) with that fixed vector n^μ as our new definition of four-momentum, the problem disappears.

Eq. (2.11) with a fixed timelike unit vector n^μ provides an alternative manifestly Lorentz-invariant definition of four-momentum. Under this new definition—which was proposed by, among others, Enrico Fermi (1922)¹⁹ and Fritz Rohrlich (1960, 1965)—the four-momentum of a spatially extended system transforms as a four-vector under Lorentz transformations no matter whether the system is open or closed. The definitions (2.9) and (2.11) are equivalent to one another for closed systems, but only coincide for open systems in the frame of reference in which n^μ has components $(1, 0, 0, 0)$. In this paper, we shall use the admittedly less elegant definition (2.9), simply because either it or its decomposition into eqs. (2.10) were the definitions used in the period of interest. Part of the problem encountered by our protagonists simply disappears by switching to the alternative definition (2.10). With this definition energy and momentum always obey the familiar relativistic transformation rules, regardless of whether we are dealing with closed systems or with their open components. As one would expect, however, a mere change of definition does not take care of the main problem that troubled the likes of Lorentz, Poincaré, and Abraham. That is the problem of the stability of a spatially extended electromagnetic electron.

¹⁸ As Gordon Fleming (private communication) has emphasized, the rest frame cannot always be uniquely defined. For the systems that will concern us here, this is not a problem. Following Fleming, one can avoid the arbitrary choice of n^μ altogether by accepting that the four-momentum of spatially extended systems is a hyperplane-dependent quantity.

¹⁹ Some of Fermi’s earliest papers are on this issue (Miller 1973, 317). We have not been able to determine what sparked Fermi’s interest in this problem. His biographer only devotes one short paragraph to it: “In January 1921, Fermi published his first paper, “On the Dynamics of a Rigid System of Electrical Charges in Translational Motion” [Fermi 1921]. This subject is of continuing interest; Fermi pursued it for a number of years and even now it occasionally appears in the literature” (Segrè 1970, 21).

2.2 Pre-relativistic theory. The analogues of relations (2.6) between energy, mass, momentum, and velocity in Newtonian mechanics are the basic formulae for kinetic energy and momentum:

$$U_{\text{kin}} = \frac{1}{2}mv^2, \quad \mathbf{p} = m\mathbf{v}. \quad (2.12)$$

In the years before the advent of special relativity, physicists worked with a hybrid theory in which Galilean-invariant Newtonian mechanics was supposed to govern matter while the inherently Lorentz-invariant electrodynamics of Maxwell and Lorentz governed the electromagnetic fields. In this hybrid theory they had already come across what are essentially the relativistic energy-momentum-velocity relations.

Initially, their starting point had still unquestionably been Newton's second law, $\mathbf{F} = m\mathbf{a}$. Electrodynamics merely supplied the Lorentz force for the left-hand side of this equation. Eventually, however, physicists were leaning toward the view that matter does not have any Newtonian mass at all and that its inertia is just a manifestation of the interaction of electric charge distributions with their self-fields. Lorentz was reluctantly driven to this conclusion because it would help explain the absence of any signs of ether drift. Abraham enthusiastically embraced it because it opened up the prospect of a purely electromagnetic basis for all of physics. With $\mathbf{F} = m\mathbf{a}$ reduced to $\mathbf{F} = 0$ Newton's second law only nominally retained its lofty position of the fundamental equation of motion. All real work was done by electrodynamics. Writing $\mathbf{F} = 0$ as $d\mathbf{P}_{\text{tot}}/dt = 0$, one can read it as expressing momentum conservation. Momentum does not need to be mechanical. Abraham introduced the concept of electromagnetic momentum.²⁰ Lorentz was happy to leave Newtonian royalty its ceremonial role. Abraham, of a more regicidal temperament, sought to replace $\mathbf{F} = m\mathbf{a}$ by a new purely electrodynamic equation that would explain why Newton's law had appeared to be the rule of the land for so long.

Despite their different motivations, Lorentz and Abraham agreed that the effective equation of motion for an electron in some external field is²¹

$$\mathbf{F}_{\text{ext}} + \mathbf{F}_{\text{self}} = 0, \quad (2.13)$$

²⁰ See (Abraham 1902a, 25–26; 1903, 110). In both places, he cites (Poincaré 1900) for the basic idea of ascribing momentum to the electromagnetic field. For discussion, see (Miller 1981, sec. 1.10), (Darrigol 1995), and (Janssen 2003, sec. 3)

²¹ In fact, another force, \mathbf{F}_{stab} , needs to be added to keep the charges from flying apart under the influence of their Coulomb repulsion.

with \mathbf{F}_{ext} the Lorentz force coming from the external field and \mathbf{F}_{self} the Lorentz force coming from the self-field of the electron. The key experiments to which eq. (2.13) was applied were the experiments of Kaufmann and others on the deflection of fast electrons by electric and magnetic fields. Both Lorentz and Abraham conceived of the electron as a spatially extended spherical surface charge distribution. They disagreed about whether the electron's shape would depend on its velocity with respect to the ether, more specifically about whether it would be subject to a microscopic version of the Lorentz-FitzGerald contraction. Lorentz believed it would, Abraham believed it would not.

The Lorentz force the electron experiences from its self-field can be written as minus the time derivative of the quantity that Abraham proposed to call the electromagnetic momentum:

$$\mathbf{F}_{\text{self}} = \int \rho(\mathbf{E} + \mathbf{v} \times \mathbf{B}) d^3x = -\frac{d\mathbf{P}_{\text{EM}}}{dt}. \quad (2.14)$$

In this expression ρ is density of the electron's charge distribution, and \mathbf{E} and \mathbf{B} are the electric and magnetic field produced by this charge distribution. The electromagnetic momentum of these fields is defined as

$$\mathbf{P}_{\text{EM}} \equiv \int \epsilon_0 \mathbf{E} \times \mathbf{B} d^3x. \quad (2.15)$$

and doubles as the electromagnetic momentum of the electron itself.

Eq. (2.14) can be derived as follows (Abraham 1905, sec. 5; Lorentz 1904a, sec. 7; Janssen 1995, 56–58). Using the inhomogeneous Maxwell equations,

$$\text{div}\mathbf{E} = \rho/\epsilon_0, \quad \text{curl}\mathbf{B} = \mu_0\rho\mathbf{v} + \frac{1}{c^2} \frac{\partial\mathbf{E}}{\partial t}, \quad (2.16)$$

we can eliminate charge and current density from the Lorentz force density, the integrand in eq. (2.14):

$$\rho(\mathbf{E} + \mathbf{v} \times \mathbf{B}) = \epsilon_0 \mathbf{E} \text{div}\mathbf{E} + \mu_0^{-1} \text{curl}\mathbf{B} \times \mathbf{B} - \frac{\mu_0^{-1}}{c^2} \frac{\partial\mathbf{E}}{\partial t} \times \mathbf{B}. \quad (2.17)$$

The last term can be written as:

$$-\frac{\partial}{\partial t}(\epsilon_0 \mathbf{E} \times \mathbf{B}) + \epsilon_0 \mathbf{E} \times \frac{\partial\mathbf{B}}{\partial t}, \quad (2.18)$$

where we used that $c^2 = 1/\epsilon_0\mu_0$. The first of these two terms is minus the time derivative of the electromagnetic momentum density (cf. eq. (2.15)). The integral over this term gives the right-hand side of eq. (2.14). A few lines of calculation show that the remaining terms in eqs. (2.17)–(2.18) combine to form the divergence of the Maxwell stress tensor. The integral over this divergence vanishes on account of Gauss' theorem. Since we shall encounter the same calculation again in sec. 6 (see eq. (6.5)), we shall go through it here in some detail.

Using the homogeneous Maxwell equations,

$$\operatorname{div}\mathbf{B} = 0, \quad \operatorname{curl}\mathbf{E} = -\frac{\partial\mathbf{B}}{\partial t}, \quad (2.19)$$

we can write the remaining terms in eqs. (2.17)–(2.18) in a form symmetric in \mathbf{E} and \mathbf{B} :

$$\epsilon_0(\mathbf{E}\operatorname{div}\mathbf{E} + \operatorname{curl}\mathbf{E} \times \mathbf{E}) + \mu_0^{-1}(\mathbf{B}\operatorname{div}\mathbf{B} + \operatorname{curl}\mathbf{B} \times \mathbf{B}). \quad (2.20)$$

The further manipulation of this expression is best done in terms of its components. We introduce $\{E^i\}_{i=1,2,3}$ and $\{B^i\}_{i=1,2,3}$ for $\{E_x, E_y, E_z\}$ and $\{B_x, B_y, B_z\}$, respectively.

The part of expression (2.20) depending on \mathbf{E} has components:

$$\epsilon_0\left(E^i\partial_k E^k + \epsilon_{ijk}\left(\epsilon_{jlm}\partial_l E^m\right)E^k\right), \quad (2.21)$$

where ϵ_{ijk} is the fully anti-symmetric Levi-Civita tensor.²² Inserting

$$\epsilon_{ijk}\epsilon_{jlm} = -\epsilon_{jik}\epsilon_{jlm} = -\delta_{il}\delta_{km} + \delta_{im}\delta_{kl}, \quad (2.22)$$

where δ_{ij} is the Kronecker delta,²³ into this expression, we find

$$\epsilon_0\left(E^i\partial_k E^k - \partial_i E^k E^k + \partial_k E^i E^k\right) = \epsilon_0\partial_k\left(E^i E^k - \frac{1}{2}\delta^{ik} E^2\right). \quad (2.23)$$

The part of eq. (2.20) depending on \mathbf{B} can likewise be written as

$$\mu_0^{-1}\partial_k\left(B^i B^k - \frac{1}{2}\delta^{ik} B^2\right). \quad (2.24)$$

²² Its definition is: $\epsilon_{ijk} = 1$ for even permutations of 123, -1 for odd permutations, and 0 otherwise.

²³ Its definition is: $\delta_{ij} = \delta^{ij} = 1$ for $i=j$ and 0 otherwise.

The sum of these two parts is the divergence of the Maxwell stress tensor,

$$T_{\text{Maxwell}}^{ij} \equiv \epsilon_0 \left(E^i E^j - \frac{1}{2} \delta^{ij} E^2 \right) + \mu_0^{-1} \left(B^i B^j - \frac{1}{2} \delta^{ij} B^2 \right). \quad (2.25)$$

Gauss' theorem tells us that

$$\int \partial_j T_{\text{Maxwell}}^{ij} d^3x = 0 \quad (2.26)$$

as long as T_{Maxwell}^{ij} drops off faster than $1/r^2$ as \mathbf{x} goes to infinity. This concludes the proof that the only term that contributes to the Lorentz force density in eq. (2.17) is the first term in eq. (2.18) with the electromagnetic momentum density.

With the help of eq. (2.14) the electromagnetic equation of motion (2.13) can be written in the form of the Newtonian equation $\mathbf{F} = d\mathbf{p}/dt$ with Abraham's electromagnetic momentum replacing ordinary momentum:

$$\mathbf{F}_{\text{ext}} = \frac{d\mathbf{P}_{\text{EM}}}{dt} \quad (2.27)$$

Like Newton's second law, which can be written either as $\mathbf{F} = m\mathbf{a}$ or as $\mathbf{F} = d\mathbf{p}/dt$, this new law can, under special circumstances, be written as the product of mass and acceleration. Assume that the momentum is in the direction of motion,²⁴ i.e., that $\mathbf{P}_{\text{EM}} = (P_{\text{EM}}/v)\mathbf{v}$. We then have

²⁴ This assumption may sound innocuous, but under the standard definition (2.9) of the four-momentum of spatially extended systems, the (ordinary three-)momentum of open systems will in general not be in the direction of motion. Because both Lorentz's and Abraham's electrons are symmetric around an axis in the direction of motion, the momentum of their self-fields is always in the direction of motion, even though these fields by themselves do not constitute closed systems. If a system has momentum that is not in the direction of motion, it will be subject to a turning couple trying to align its momentum with its velocity. Trouton and Noble (1903) tried in vain to detect this effect on a charged capacitor hanging from the ceiling of their laboratory on a torsion wire (Janssen 2002b, 440–441, note; Janssen 1995, especially secs. 1.4.2 and 2.2.5). Ehrenfest (1907) raised the question whether the electron would be subject to a turning couple if it were *not* symmetric around the axis in the direction of motion. Einstein (1907a) countered that the behavior of the electron would be independent of its shape (see Miller 1981, sec. 7.4.4, for discussion of this exchange between Einstein and Ehrenfest). Laue (1911a) showed how this could be (see also Pauli 1921, 186–187). As with the capacitor in the Trouton-Noble experiment, the electromagnetic momentum of the electron is not the only momentum of the system. The non-electromagnetic part of the system also contributes to its momentum. Laue showed that the total momentum of a closed static system is always in the direction of motion. From a modern point of view this is a direct consequence of the fact that four-

$$\frac{d\mathbf{P}_{\text{EM}}}{dt} = \frac{dP_{\text{EM}}}{dt} \frac{\mathbf{v}}{v} + P_{\text{EM}} \frac{d}{dt} \left(\frac{\mathbf{v}}{v} \right). \quad (2.28)$$

The first term on the right-hand side can be written as

$$\frac{dP_{\text{EM}}}{dt} \frac{\mathbf{v}}{v} = \frac{dP_{\text{EM}}}{dv} \frac{dv}{dt} \frac{\mathbf{v}}{v} = \frac{dP_{\text{EM}}}{dv} \mathbf{a}_{\parallel}, \quad (2.29)$$

where \mathbf{a}_{\parallel} is the longitudinal acceleration, i.e., the acceleration in the direction of motion.

The second term can be written as

$$P_{\text{EM}} \frac{d}{dt} \left(\frac{\mathbf{v}}{v} \right) = \frac{P_{\text{EM}}}{v} \mathbf{a}_{\perp}, \quad (2.30)$$

where \mathbf{a}_{\perp} is the transverse acceleration, i.e., the acceleration perpendicular to the direction of motion. The factors multiplying these two components of the acceleration are called the *longitudinal mass*, m_{\parallel} , and the *transverse mass*, m_{\perp} , respectively. Eq. (2.28) can thus be written as

$$\frac{d\mathbf{P}_{\text{EM}}}{dt} = m_{\parallel} \mathbf{a}_{\parallel} + m_{\perp} \mathbf{a}_{\perp}, \quad (2.31)$$

with²⁵

$$m_{\parallel} = \frac{dP_{\text{EM}}}{dv}, \quad m_{\perp} = \frac{P_{\text{EM}}}{v}. \quad (2.32)$$

The effective equation of motion (2.27) becomes:

$$\mathbf{F}_{\text{ext}} = m_{\parallel} \mathbf{a}_{\parallel} + m_{\perp} \mathbf{a}_{\perp}, \quad (2.33)$$

momentum of a closed system (static or not) transforms as a four-vector under Lorentz transformations. The momenta of open systems, such as the subsystems of a closed static system, need not be in the direction of motion, in which case the system is subject to equal and opposite turning couples. A closed system never experiences a net turning couple. The turning couples on open systems, it turns out, are artifacts of the standard definition (2.9) of the four-momentum of spatially extended systems. Under the alternative Fermi-Rohrlich definition (see the discussion following eq. (2.11)), there are no turning couples whatsoever (Butler 1968, Janssen 1995, Teukolsky 1996).

²⁵ Substituting the momentum, $p=mv$, of Newtonian mechanics for P_{EM} in eq. (2.32), we find $m_{\parallel}=m_{\perp}=m$.

We shall see that for $v = 0$ (in which case the electron models of Abraham and Lorentz coincide) $m_{//} = m_{\perp} = m_0$, and that for $v \neq 0$ $m_{//}$ and m_{\perp} differ from m_0 only by terms of order v^2/c^2 . For velocities $v \ll c$, eq. (2.33) thus reduces to:

$$\mathbf{F}_{\text{ext}} \approx m_0(\mathbf{a}_{//} + \mathbf{a}_{\perp}) = m_0\mathbf{a} \quad (2.34)$$

Proponents of the electromagnetic view of nature took eq. (2.27) to be the fundamental equation of motion and derived Newton's law from it by identifying the ordinary Newtonian mass with the electromagnetic mass m_0 of the relevant system at rest in the ether.

Eq. (2.32) defines the longitudinal mass $m_{//}$ of the electron in terms of its electromagnetic momentum. It can also be defined in terms of the electron's electromagnetic energy. Consider the work done as an electron is moving in the x -direction in the absence of an external field. The work expended goes into the internal energy of the electron, $dU = -dW$. According to eq. (2.13), the work is done by \mathbf{F}_{self} .²⁶ The internal energy is identified with the electromagnetic energy U_{EM} .

$$dU_{\text{EM}} = -dW = -\mathbf{F}_{\text{self}} \cdot d\mathbf{x}. \quad (2.35)$$

Using eqs. (2.14) and (2.31), we can write this as

$$dU_{\text{EM}} = \frac{d\mathbf{P}_{\text{EM}}}{dt} \cdot d\mathbf{x} = m_{//}\mathbf{a}_{//} \cdot d\mathbf{x} = m_{//} \frac{dv}{dt} dx = m_{//}v dv \quad (2.36)$$

It follows that²⁷

$$m_{//} = \frac{1}{v} \frac{dU_{\text{EM}}}{dv} \quad (2.37)$$

As we shall see in sec. 4, given the standard definitions (2.10) of electromagnetic energy and momentum, the neglect of non-electromagnetic stabilizing forces in the derivation of eqs. (2.32) and (2.37) leads to an ambiguity in the expression for the longitudinal mass of Lorentz's electron.

²⁶ But recall that there should be an additional term, \mathbf{F}_{stab} , on the right-hand side of eq. (2.13) (see note 21).

²⁷ Substituting the kinetic energy, $U_{\text{kin}} = \frac{1}{2}mv^2$, of Newtonian mechanics for U_{EM} in eq. (2.37), we find $m_{//} = m$, in accordance with the result found on the basis of eq. (2.32) and $p=mv$ (see note 25).

If the combination of the energy U (divided by c), and the momentum \mathbf{P} for any system, electromagnetic or otherwise, transforms as a four-vector under Lorentz transformations, then $m_{//}$ calculated from eq. (2.32) (with P substituted for P_{EM}) is equal to $m_{//}$ calculated from eq. (2.37) (with U substituted for U_{EM}).²⁸ Consider the transformation from a rest frame with coordinates x_0^μ to the x^μ -frame. In that case (see eqs. (2.1)–(2.5)):

$$P^\mu = \left(\frac{U}{c}, \mathbf{P} \right) = (\gamma m_0 c, \gamma m_0 \mathbf{v}). \quad (2.38)$$

The energy U gives the longitudinal mass (see eq. (2.37))

$$m_{//} = \frac{1}{v} \frac{dU}{dv} = \frac{1}{v} \frac{d}{dv} (\gamma m_0 c^2) = \frac{m_0 c^2}{v} \frac{d\gamma}{dv}. \quad (2.39)$$

The momentum \mathbf{P} gives the longitudinal mass (eq. (2.32)):

$$m_{//} = \frac{dP}{dv} = \frac{d}{dv} (\gamma m_0 v) = m_0 \frac{d(\gamma v)}{dv}, \quad (2.40)$$

Using that^{29, 30}

$$\frac{d\gamma}{dv} = \gamma^3 \frac{v}{c^2}, \quad \frac{d(\gamma v)}{dv} = \gamma^3, \quad (2.41)$$

we find that eqs. (2.39) and (2.40) do indeed give the same result:

$$m_{//} = \frac{1}{v} \frac{dU}{dv} = \frac{dP}{dv} = \gamma^3 m_0, \quad (2.42)$$

The momentum \mathbf{P} in eq. (2.38) gives the transverse mass (eq. (2.32)):

²⁸ The converse is not true. For the electron model of Bucherer and Langevin (see sec. 4) ($U_{\text{EM}}/c, \mathbf{P}_{\text{EM}}$) is not a four-vector, yet U_{EM} and \mathbf{P}_{EM} give the same longitudinal mass $m_{//}$ (see eqs. (4.23)–(4.26)). The same is true for the Newtonian energy $U_{\text{kin}} = \frac{1}{2} m v^2$ and the Newtonian momentum $p = m v$ (see notes 25 and 27).

²⁹ $\frac{d\gamma}{dv} = \frac{d\gamma}{d\beta} \frac{d\beta}{dv} = \frac{d}{d\beta} (1-\beta^2)^{-1/2} \frac{1}{c} = -\frac{1}{2} (1-\beta^2)^{-3/2} \frac{d}{d\beta} (1-\beta^2) \frac{1}{c} = -\frac{1}{2} \gamma^3 (-2\beta) \frac{1}{c} = \gamma^3 \frac{v}{c^2}$.

³⁰ $\frac{d(\gamma v)}{dv} = \gamma + v \frac{d\gamma}{dv} = \gamma + \gamma^3 \beta^2 = \gamma^3 (1-\beta^2 + \beta^2) = \gamma^3$

$$m_{\perp} = \frac{P}{v} = \gamma m_0 \quad (2.43)$$

Eqs. (2.42) and (2.43) give mass-velocity relations that hold for any relativistic particle. These equations thus have much broader applicability than their origin in electrodynamics suggests. This is exactly what killed the dreams of Abraham and Lorentz of using these relations to draw conclusions about the nature and shape of the electron.

3. Lorentz's theorem of corresponding states, the generalized contraction hypothesis, and the velocity dependence of electron mass

Lorentz had already published the relativistic eqs. (2.42) and (2.43) for longitudinal and transverse mass, up to an undetermined factor l , in 1899. To understand how Lorentz originally arrived at these equations we need to take a look at his general approach to problems in the electrodynamics of moving bodies.³¹ The basic problem that Lorentz was facing was that Maxwell's equations are not invariant under Galilean transformations, which relate frames in relative motion to one another in Lorentz's classical Newtonian space-time. Lorentz thus labored under the impression that Maxwell's equations only hold in frames at rest in the ether and not in the terrestrial lab frames in which all our experiments are done.

Consider an ether frame with space-time coordinates (t_0, \mathbf{x}_0) and a lab frame with space-time coordinates (t, \mathbf{x}) related to one another via the Galilean transformation

$$t = t_0, \quad x = x_0 - vt_0, \quad y = y_0, \quad z = z_0, \quad (3.1)$$

$$\mathbf{E} = \mathbf{E}_0, \quad \mathbf{B} = \mathbf{B}_0, \quad \rho = \rho_0$$

The second line of this equation expresses that the electric field, the magnetic field, and the charge density remain the same even though after the transformation they are thought of as functions of (t, \mathbf{x}) rather than as functions of (t_0, \mathbf{x}_0) .

The equations for the fields produced by a charge distribution static in the lab frame as functions of the space-time coordinates (t, \mathbf{x}) are obtained by writing down Maxwell's equations for the relevant quantities in the lab frame, adding the current $\mu_0 \rho \mathbf{v}$ ³² and replacing time derivatives by the operator $\partial/\partial t - v\partial/\partial x$.³³ We thus arrive at:

$$\begin{aligned} \operatorname{div} \mathbf{E} &= \rho / \epsilon_0 & \operatorname{curl} \mathbf{B} &= \mu_0 \rho \mathbf{v} + \frac{1}{c^2} \left(\frac{\partial \mathbf{E}}{\partial t} - v \frac{\partial \mathbf{E}}{\partial x} \right) \\ \operatorname{div} \mathbf{B} &= 0 & \operatorname{curl} \mathbf{E} &= -\frac{\partial \mathbf{B}}{\partial t} + v \frac{\partial \mathbf{B}}{\partial x} \end{aligned} \quad (3.2)$$

³¹ For more extensive discussion, see (Janssen 1995, Ch. 3; 2002b; Janssen and Stachel 2004).

³² For the magnetic field it is the motion of charges with respect to the ether that matters, not the motion with respect to the lab frame.

³³ For the induced \mathbf{E} and \mathbf{B} fields it is the changes in the \mathbf{B} and \mathbf{E} fields at fixed points in the ether that matter, not the changes at fixed points in the lab frame.

Lorentz now replaced the space-time coordinates (t, \mathbf{x}) , the fields \mathbf{E} and \mathbf{B} , and the charge density ρ by auxiliary variables defined as:

$$\begin{aligned} \mathbf{x}' &= l \operatorname{diag}(\gamma, 1, 1) \mathbf{x}, \quad t' = l \left(\frac{t}{\gamma} - \gamma \left(\frac{\mathbf{v}}{c^2} \right) \cdot \mathbf{x} \right), \quad \rho' = \frac{\rho}{\gamma l^3}, \\ \mathbf{E}' &= \frac{1}{l^2} \operatorname{diag}(1, \gamma, \gamma) (\mathbf{E} + \mathbf{v} \times \mathbf{B}), \quad \mathbf{B}' = \frac{1}{l^2} \operatorname{diag}(1, \gamma, \gamma) \left(\mathbf{B} - \frac{1}{c^2} \mathbf{v} \times \mathbf{E} \right), \end{aligned} \quad (3.3)$$

where l is an undetermined factor that is assumed to be equal to one to first order in v/c . Since the auxiliary time variable depends on position, it is called local time. He showed that the auxiliary fields \mathbf{E}' and \mathbf{B}' and the auxiliary charge density ρ' written as functions of the auxiliary space-time coordinates (t', \mathbf{x}') satisfy Maxwell's equations:

$$\begin{aligned} \operatorname{div}' \mathbf{E}' &= \rho' / \epsilon_0 & \operatorname{curl}' \mathbf{B}' &= \frac{1}{c^2} \frac{\partial \mathbf{E}'}{\partial t'} \\ \operatorname{div}' \mathbf{B}' &= 0 & \operatorname{curl}' \mathbf{E}' &= - \frac{\partial \mathbf{B}'}{\partial t'} \end{aligned} \quad (3.4)$$

When the factor l is set exactly equal to one, what Lorentz showed, at least for static charge densities,³⁴ is that Maxwell's equations are invariant under what Poincaré (1906, 495) proposed to call *Lorentz transformations*. For $l = 1$, the transformation formulae in eq. (3.3) for the fields \mathbf{E} and \mathbf{B} and for a static charge density ρ look exactly the same as in special relativity. The transformation formulae for the space-time coordinates do not. Bear in mind, however, that Lorentz did the transformation in two steps, given by eqs. (3.1) and (3.3), respectively. Schematically, we have:

$$(t_0, \mathbf{x}_0, \mathbf{E}_0, \mathbf{B}_0, \rho_0) \rightarrow (t, \mathbf{x}, \mathbf{E}, \mathbf{B}, \rho) \rightarrow (t', \mathbf{x}', \mathbf{E}', \mathbf{B}', \rho')_{l=1} \quad (3.5)$$

Combining these two steps, we recover the familiar Lorentz transformation formulae. For the fields and the charge density, this is just a matter of replacing $(\mathbf{E}, \mathbf{B}, \rho)$ in eq. (3.3) by $(\mathbf{E}_0, \mathbf{B}_0, \rho_0)$. For the space-time coordinates, it takes only a minimal amount of algebra:

³⁴ Lorentz only started using the relativistic transformation formula for non-static charge densities and for current densities in 1915 (see Janssen 1995, secs. 3.5.3 and 3.5.6).

$$x' = \gamma x = \gamma(x_0 - vt_0), \quad y' = y = y_0, \quad z' = z = z_0 \quad (3.6)$$

$$t' = \frac{t}{\gamma} - \gamma \frac{v}{c^2} x = \frac{t_0}{\gamma} - \gamma \frac{v}{c^2} (x_0 - vt_0) = \gamma \left(t_0 - \frac{v}{c^2} x_0 \right)$$

where in the second line we used that $1/\gamma + \gamma(v^2/c^2) = \gamma(1/\gamma^2 + \beta^2) = \gamma$.

The inverse of the transformation $(t_0, \mathbf{x}_0, \mathbf{E}_0, \mathbf{B}_0) \rightarrow (t', \mathbf{x}', \mathbf{E}', \mathbf{B}')$ for $l=1$ is found by interchanging $(t_0, \mathbf{x}_0, \mathbf{E}_0, \mathbf{B}_0)$ and $(t', \mathbf{x}', \mathbf{E}', \mathbf{B}')$ and changing \mathbf{v} to $-\mathbf{v}$. Doing the inversion for $l \neq 1$ also requires changing l to l^{-1} . The inverse of the transformation $(\mathbf{E}_0, \mathbf{B}_0) \rightarrow (\mathbf{E}', \mathbf{B}')$ for $l \neq 1$, for instance, is given by

$$\begin{aligned} \mathbf{E}_0 = \mathbf{E} &= l^2 \text{diag}(1, \gamma, \gamma) (\mathbf{E}' - \mathbf{v} \times \mathbf{B}') \\ \mathbf{B}_0 = \mathbf{B} &= l^2 \text{diag}(1, \gamma, \gamma) \left(\mathbf{B}' + \frac{1}{c^2} \mathbf{v} \times \mathbf{E}' \right) \end{aligned} \quad (3.7)$$

The transformation is symmetric only for $l=1$. Unlike Lorentz before 1905, Poincaré and Einstein both looked upon the primed quantities as the quantities measured by the observer in the lab frame. In special relativity, the ether frame is just another inertial frame on a par with the lab frame. The situation for observers in these two frames will be fully symmetric only if $l=1$. This was essentially the argument for both Poincaré and Einstein to set $l=1$. As we shall see in the next section, Lorentz also ended up setting $l=1$ but on the basis of a roundabout dynamical argument. For our purposes it is important that we leave the factor l undetermined for the time being.

The invariance of Maxwell's equations under the combination of transformations (3.1) and (3.3) allowed Lorentz to formulate what he called the *theorem of corresponding states*. This theorem says that for any field configuration in a frame at rest in the ether there is a corresponding field configuration in a frame moving through the ether such that the auxiliary fields \mathbf{E}' and \mathbf{B}' in the moving frame are the same functions of the auxiliary space and time coordinates (t', \mathbf{x}') as the real fields \mathbf{E}_0 and \mathbf{B}_0 in the frame at rest of the real space and time coordinates (t_0, \mathbf{x}_0) . Lorentz was particularly interested in free field configurations (for which $\rho = 0$) describing patterns of light and darkness. Most experiments in optics eventually boil down to the observation of such patterns.

To describe a pattern of light and darkness it suffices to specify where the fields averaged over times that are long compared to the period of the light used vanish and where these averages are large. \mathbf{E}' and \mathbf{B}' are linear combinations of \mathbf{E} and \mathbf{B} (see eq.

(3.3)). They are large (small) when- and wherever \mathbf{E} and \mathbf{B} are. Since patterns of light and darkness by their very nature are effectively static, no complications arise from the x -dependence of local time. If it is light (dark) simultaneously at two points with coordinates $\mathbf{x}_0 = \mathbf{a}$ and $\mathbf{x}_0 = \mathbf{b}$ in some field configuration in a frame at rest in the ether, it will be light (dark) simultaneously at the corresponding points $\mathbf{x}' = \mathbf{a}$ and $\mathbf{x}' = \mathbf{b}$ in the corresponding state in a frame moving through the ether. In terms of the real coordinates these are the points $\mathbf{x} = (1/l)\text{diag}(1/\gamma, 1, 1)\mathbf{a}$ and $\mathbf{x} = (1/l)\text{diag}(1/\gamma, 1, 1)\mathbf{b}$. The pattern of light and darkness in a moving frame is thus obtained from its corresponding pattern in a frame at rest in the ether by contracting the latter by a factor γl in the direction of motion and a factor l in the directions perpendicular to the direction of motion. Examining the formula for the local time in eq. (3.3), one likewise sees that the periods of light waves in a moving frame are obtained by multiplying the periods of the light waves in the corresponding state at rest in the ether by a factor γ/l .

To account for the fact that these length-contraction and time-dilation effects in electromagnetic field configurations were never detected, Lorentz (1899) assumed that matter interacting with the fields (e.g., the optical components producing patterns of light and darkness) experiences these same effects. Lorentz thereby added a far-reaching physical assumption to his purely mathematical theorem of corresponding states. Elsewhere one of us has dubbed this assumption the *generalized contraction hypothesis* (Janssen 1995, sec. 3.3; 2002b; Janssen and Stachel 2004). It was through this hypothesis that Lorentz decreed a number of exceptions to the Newtonian laws that had jurisdiction over matter in his theory. The length-contraction and time-dilation rules to which matter and field alike had to be subject to account for the absence of any signs of ether drift are examples of such exceptions. The velocity dependence of mass is another (Janssen 1995, sec. 3.3.6). It is this exception that is important for our purposes.

Consider an oscillating electron generating an electromagnetic wave in a light source in the frame S_0 at rest in the ether. Suppose the oscillation satisfies Newton's laws of motion. Now consider that same electron in the corresponding state in the Galilean frame S moving through the ether with velocity \mathbf{v} . In terms of the auxiliary space-time coordinates in eq. (3.3) its motion will be exactly the same as the motion in the system at rest. This implies that in terms of the real quantities, it will not satisfy Newton's laws of motion, unless the electron's mass depends on its velocity in a specific way. Lorentz (1899) set out to determine that velocity dependence.

Suppose an oscillating electron in a light source at rest in S_0 satisfies $\mathbf{F}_0 = m_0\mathbf{a}_0$. In the corresponding state in S the corresponding electron will then satisfy the same equation in terms of the auxiliary quantities, i.e.,

$$\mathbf{F}' = m_0 \mathbf{a}' , \quad (3.8)$$

where \mathbf{F}' is the same function of (t', \mathbf{x}') as \mathbf{F}_0 is of (t_0, \mathbf{x}_0) , and where $\mathbf{a}' = d^2 \mathbf{x}' / dt'^2$ and $\mathbf{a}_0 = d^2 \mathbf{x}_0 / dt_0^2$ are always the same at corresponding points in S and S_0 . Lorentz assumed that motion through the ether affects all forces on the electron the same way it affects Coulomb forces³⁵

$$\mathbf{F}' = \frac{1}{l^2} \text{diag}(1, \gamma, \gamma) \mathbf{F} . \quad (3.9)$$

For the relation between the acceleration \mathbf{a}' in terms of the auxiliary space and time coordinates and the real acceleration \mathbf{a} , Lorentz used the relation

$$\mathbf{a}' = \frac{1}{l} \text{diag}(\gamma^3, \gamma^2, \gamma^2) \mathbf{a} . \quad (3.10)$$

In general, this relation is far more complicated, but when the velocity $d\mathbf{x}_0 / dt_0$ with which the electron is oscillating in S_0 is small, $d\mathbf{x}' / dt'$ (equal to $d\mathbf{x}_0 / dt_0$ at the corresponding point in S_0) can be neglected and eq. (3.10) holds. A derivation of the general relation between \mathbf{a}' and \mathbf{a} was given by Planck (1906a) in the context of his derivation of the relativistic generalization of Newton's second law, a derivation mathematically essentially equivalent to Lorentz's 1899 derivation of the velocity dependence of mass, except that Planck only had to consider the special case $l = 1$.³⁶

Lorentz probably arrived at eq. (3.10) through the following crude argument. If an electron oscillates around a fixed point in S with a low velocity and a small amplitude, the x -dependent term in the expression for local time can be ignored. In that case, we only need to take into account that \mathbf{x}' differs from \mathbf{x} by a factor $l \text{diag}(\gamma, 1, 1)$ and that t' differs from t by a factor l/γ (see eq. (3.3)). This gives a quick and dirty derivation of eq. (3.10):

$$\mathbf{a}' = \frac{d^2 \mathbf{x}'}{dt'^2} = \left(\frac{\gamma}{l}\right)^2 l \text{diag}(\gamma, 1, 1) \frac{d^2 \mathbf{x}}{dt^2} = \frac{1}{l} \text{diag}(\gamma^3, \gamma^2, \gamma^2) \mathbf{a} . \quad (3.11)$$

Inserting eqs. (3.9) and (3.10) into eq. (3.8), we find

³⁵ See (Lorentz 1895, sec. 19–23) for the derivation of this transformation law and (Janssen 1995, sec. 3.2.5) or (Zahar 1989, 59–61) for a reconstruction of the derivation in modern notation.

³⁶ For an elegant and elementary exposition of Planck's derivation, see (Zahar 1989, sec. 7.1, 227–237). The equations for the relation between \mathbf{a}' and \mathbf{a} can be found on p. 232, eqs. (2)–(4).

$$\frac{1}{l^2} \text{diag}(1, \gamma, \gamma) \mathbf{F} = \frac{1}{l} \text{diag}(\gamma^3, \gamma^2, \gamma^2) m_0 \mathbf{a} \quad (3.12)$$

This can be rewritten as

$$\mathbf{F} = l \text{diag}(\gamma^3, \gamma, \gamma) m_0 \mathbf{a}. \quad (3.13)$$

From this equation it follows that the oscillation of an electron in the moving source can only satisfy Newton's second law if the mass m of an electron with velocity \mathbf{v} with respect to the ether (remember that the velocity of the oscillation itself was assumed to be negligible) differs from the mass m_0 of an electron at rest in the ether in precisely the following way:

$$m_{//} = l\gamma^3 m_0, \quad m_{\perp} = l\gamma m_0. \quad (3.14)$$

If $l = 1$, these are just the relativistic eqs. (2.42) and (2.43). It was Planck who showed in the paper mentioned above that these relations also obtain in special relativity.³⁷ Planck's interpretation of these relations was very different from Lorentz's. For Planck, as for Einstein, the velocity dependence of mass was part of a new relativistic mechanics replacing classical Newtonian mechanics. Lorentz wanted to retain Newtonian mechanics, even after he accepted in 1904 that there are no Galilean-invariant Newtonian masses or forces in nature. Consequently, he had to provide an explanation for the peculiar velocity-dependence of electron mass he needed to account for the absence of any detectable ether drift. In 1904, adapting Abraham's electron model, Lorentz provided such an explanation in the form of a specific model of the electron that exhibited exactly the velocity dependence of eq. (3.14) for $l = 1$.

³⁷ Einstein (1905, 919) obtained $m_{\perp} = \gamma^2 m_0$ instead of $m_{\perp} = \gamma m_0$, the result obtained by Planck and Lorentz (for $l=1$). The discrepancy comes from Einstein using $\mathbf{F}' = \mathbf{F}$ instead of $\mathbf{F}' = \text{diag}(1, \gamma, \gamma) \mathbf{F}$, the now standard transformation law for forces used by Lorentz and Planck (Zahar 1989, p. 233). Einstein made it clear that he was well aware of the arbitrariness of his definition of force.

4. Electromagnetic energy, momentum, and mass of a moving electron

In this section we use Lorentz's theorem of corresponding states to calculate the energy, the momentum, and the Lagrangian for the field of a moving electron, conceived of as nothing but a surface charge distribution and its electromagnetic field. We then compute the longitudinal and the transverse mass of the electron.







| | moving electron | corresponding state stretch dimensions of moving system by $\text{diag}(\gamma l, l, l)$ |
|---|--|--|
| The rigid electron of Abraham $(l = 1)$ |  sphere (R, R, R) |  ellipsoid $(\gamma R, R, R)$ |
| The contractile electron of Lorentz and Poincaré $(l = 1)$ $\left(\frac{R}{\gamma l}, \frac{R}{l}, \frac{R}{l}\right) = \left(\frac{R}{\gamma}, R, R\right)$ |  ellipsoid $(R/\gamma, R, R)$ |  sphere (R, R, R) |
| The contractile electron of constant volume of Bucherer and Langevin $(l = \gamma^{-1/3})$ $\left(\frac{R}{\gamma l}, \frac{R}{l}, \frac{R}{l}\right) =$ $\left(\frac{R}{\gamma^{2/3}}, \gamma^{1/3}R, \gamma^{1/3}R\right)$ |  ellipsoid $R/\gamma^{2/3}, \gamma^{1/3}R, \gamma^{1/3}R$ |  sphere (R, R, R) |

Figure 1: A moving electron according to the models of Abraham, Lorentz, and Bucherer-Langevin, and the corresponding states at rest in the ether.

We distinguish three different models. In all three the electron at rest in the ether is spherical. In Abraham's model it remains spherical when it is set in motion; in Lorentz's model it contracts by a factor γ in the direction of motion; and in the Bucherer-Langevin model it contracts by a factor $\gamma^{2/3}$ in the direction of motion but expands by a factor $\gamma^{1/3}$ in the directions perpendicular to the direction of motion so that its volume remains constant. Fig. 1 shows a moving electron according to these three models along with the corresponding states in a frame at rest in the ether. For Abraham's rigid electron the corresponding state is an ellipsoid; for the contractile electrons of Lorentz and Bucherer-Langevin it is a sphere.

In the corresponding state of a moving electron (in relativistic terms: in the electron's rest frame) there is no magnetic field. Hence $\mathbf{B}' = 0$ and eq. (3.7) gives:

$$\mathbf{E} = l^2(E'_x, \gamma E'_y, \gamma E'_z), \quad \mathbf{B} = \frac{\gamma l^2 v}{c^2}(0, -E'_z, E'_y). \quad (4.1)$$

4.1 Energy. The energy of the electric and magnetic field is defined as

$$U_{\text{EM}} = \int \left(\frac{1}{2} \epsilon_0 E^2 + \frac{1}{2} \mu_0^{-1} B^2 \right) d^3 x. \quad (4.2)$$

For the field in eq. (4.1) with $B_x = 0$, it is given by

$$U_{\text{EM}} = \int \frac{1}{2} \epsilon_0 E_x^2 d^3 x + \int \frac{1}{2} \epsilon_0 (E_y^2 + E_z^2) d^3 x + \int \frac{1}{2} \mu_0^{-1} (B_y^2 + B_z^2) d^3 x. \quad (4.3)$$

Following Poincaré (1906, 523), we call these three terms A , B , and C . Using eq. (4.1) and $d^3 x = d^3 x' / \gamma l^3$, we find

$$\begin{aligned} A &= \frac{l}{\gamma} \int \frac{1}{2} \epsilon_0 E_x'^2 d^3 x' = \frac{l}{\gamma} A', \\ B &= l\gamma \int \frac{1}{2} \epsilon_0 (E_y'^2 + E_z'^2) d^3 x' = l\gamma B', \\ C &= \frac{\mu_0^{-1} \gamma l v^2}{c^4} \int \frac{1}{2} (E_y'^2 + E_z'^2) d^3 x' = l\gamma \beta^2 B'. \end{aligned} \quad (4.4)$$

If the corresponding state is spherical,

$$B' = 2A' = \frac{2}{3} U'_{\text{EM}}. \quad (4.5)$$

It follows that for the models of Lorentz and Bucherer-Langevin:

$$U_{\text{EM}} = l\gamma \left(\frac{1}{\gamma^2} + 2 + 2\beta^2 \right) A' = l\gamma \left(1 + \frac{1}{3}\beta^2 \right) U'_{\text{EM}}, \quad (4.6)$$

where we used that $\gamma^{-2} = 1 - \beta^2$ and that $3A' = U'_{\text{EM}}$. Eq. (4.6) can also be written as

$$U_{\text{EM}} = l\gamma \left(\frac{4}{3} - \frac{1}{3}(1 - \beta^2) \right) U'_{\text{EM}} = l \left(\frac{4\gamma}{3} - \frac{1}{3\gamma} \right) U'_{\text{EM}}. \quad (4.7)$$

4.2 Lagrangian. The Lagrangian can be computed the same way. We start from

$$L_{\text{EM}} = \int \mathcal{L}_{\text{EM}} d^3x, \quad (4.8)$$

where \mathcal{L}_{EM} is the Lagrange density defined as (note the sign)

$$\mathcal{L}_{\text{EM}} \equiv \frac{1}{2} \mu_0^{-1} B^2 - \frac{1}{2} \varepsilon_0 E^2. \quad (4.9)$$

This quantity transforms as a scalar under Lorentz transformations as can be seen from its definition in manifestly Lorentz-invariant form:³⁸

$$\mathcal{L}_{\text{EM}} \equiv \frac{1}{4} \mu_0^{-1} F_{\mu\nu} F^{\mu\nu}. \quad (4.10)$$

It follows that $\mathcal{L}_{\text{EM}} = l^4 \mathcal{L}'_{\text{EM}}$ with $\mathcal{L}'_{\text{EM}} = -(1/2)\varepsilon_0 E'^2$

$$L_{\text{EM}} = \int l^4 \mathcal{L}'_{\text{EM}} \frac{d^3x'}{\gamma l^3} = -\frac{l}{\gamma} U'_{\text{EM}}. \quad (4.11)$$

4.3 Momentum. The electromagnetic momentum can also be computed in this way. For the field of the electron, the electromagnetic momentum density (see eq. (2.15)) is:

$$\mathbf{p}_{\text{EM}} = \varepsilon_0 \begin{pmatrix} E_y B_z - E_z B_y \\ -E_x B_z \\ E_x B_y \end{pmatrix} = \varepsilon_0 \gamma l^4 \frac{v}{c^2} \begin{pmatrix} \gamma (E_y'^2 + E_z'^2) \\ E'_x E'_y \\ E'_x E'_z \end{pmatrix}. \quad (4.12)$$

³⁸ See eq. (6.2) below for the relation between the (contravariant) electromagnetic field strength tensor $F^{\mu\nu}$ (and its covariant form $F_{\mu\nu} = \eta_{\mu\rho} \eta_{\nu\sigma} F^{\rho\sigma}$) and the components of \mathbf{E} and \mathbf{B} .

Because of symmetry (in all three models)

$$\int p_{y_{EM}} d^3x = \int p_{z_{EM}} d^3x = 0. \quad (4.13)$$

For the x -component, we find

$$P_{x_{EM}} = \frac{1}{\gamma l^3} \int p_{x_{EM}} d^3x' = \gamma l \frac{v}{c^2} \int \epsilon_0 (E_y'^2 + E_z'^2) d^3x'. \quad (4.14)$$

This is equal to $\gamma l (v/c^2) 2B'$. For the contractile electron (Lorentz and Bucherer-Langevin), $B' = (2/3)U'_{EM}$. In that case

$$\mathbf{P}_{EM} = \frac{4}{3} \gamma l \left(\frac{U'_{EM}}{c^2} \right) \mathbf{v}. \quad (4.15)$$

This pre-relativistic equation will immediately strike anyone familiar with the basic formulae of special relativity as odd. Remember that from a relativistic point of view the energy U'_{EM} of the moving electron's corresponding state at rest in the ether is nothing but the energy U_{0EM} of the electron in its rest frame. Comparison of eq. (4.15) with $l = 1$ to $\mathbf{P} = \gamma m_0 \mathbf{v}$ (eq. (2.6)) suggests that the rest mass of the electron is $m_0 = \frac{4}{3} U_{0EM} / c^2$, which seems to be in blatant contradiction to the equation everybody knows, $E = mc^2$. This is the notorious “4/3 puzzle” of the energy-mass relation of the classical electron. The origin of the problem is that the system we are considering, the self-field of the electron, is not closed and that its four-momentum consequently does not transform as a four-vector, at least not under the standard definition (2.9). The solution to the puzzle is either to add another piece to the system so that the composite system is closed or to adopt the alternative Fermi-Rohrlich definition (2.11) (with a fixed unit vector n^μ) of the four-momentum of spatially extended systems. As we shall see, the “4/3 puzzle” had already reared its ugly head before the advent of special relativity, albeit in a different guise.

4.4 Longitudinal and transverse mass. Substituting eqs. (4.7) and (4.15) for the energy and momentum of the field of a moving contractile electron into the expressions (2.32) and (2.37) for the electron's transverse and longitudinal mass, we find:

$$m_{//} = \frac{dP_{EM}}{dv} = \frac{d(\gamma lv)}{dv} \frac{4}{3} \frac{U'_{EM}}{c^2}, \quad (4.16)$$

$$m_{\perp} = \frac{P_{EM}}{v} = \gamma l \frac{4 U'_{EM}}{3 c^2}, \quad (4.17)$$

$$m_{//} = \frac{1}{v} \frac{dU_{EM}}{dv} = \frac{1}{v} \frac{d}{dv} \left(\frac{4\gamma l}{3} - \frac{l}{3\gamma} \right) U'_{EM}. \quad (4.18)$$

Several conclusions can be drawn from these equations. First, it turns out that eq. (4.16) only gives the velocity dependence of the longitudinal mass required by Lorentz's generalized contraction hypothesis for $l = 1$. Unfortunately, for $l = 1$, eq. (4.18) does not give the same longitudinal mass as eq. (4.16). One only obtains the same result for $l = \gamma^{-1/3}$. This is the value for the Bucherer-Langevin constant-volume contractile electron.

It is easy to prove these claims. Using eq. (2.41), we can write eq. (4.16) as

$$m_{//} = \frac{dP_{EM}}{dv} = \left(\gamma^3 l + \gamma v \frac{dl}{dv} \right) \frac{4 U'_{EM}}{3 c^2}. \quad (4.19)$$

From eqs. (4.17) and (4.19) it follows that the only way to ensure that $m_{//} = l\gamma^3 m_0$ and $m_{\perp} = l\gamma m_0$, as required by the generalized contraction hypothesis (see eq. (3.14)), is to set the Newton mass equal to zero, to set $l = 1$, and to define the mass of the electron at rest in the ether as

$$m_0 = \frac{4 U'_{EM}}{3 c^2} \quad (4.20)$$

(which, from a relativistic point of view, amounts to the odd equation $E = \frac{3}{4} mc^2$). Eqs. (4.19) and (4.17) then reduce to

$$m_{//} = \gamma^3 m_0, \quad m_{\perp} = \gamma m_0, \quad (4.21)$$

in accordance with eq. (3.14).

Lorentz (1904) had thus found a concrete model for the electron with a mass exhibiting exactly the velocity dependence that he had found in 1899. This could hardly be a coincidence. Lorentz concluded³⁹ that the electron was indeed nothing but a small

³⁹ This is an example of what I called a "common origin inference" or *COI* in (Janssen 2002a). The example illustrates how easy it is to overreach with this kind of argument (for other examples see, *ibid.*, 474, 491, 508).

spherical surface charge distribution, subject to a microscopic version of the Lorentz-FitzGerald contraction when set in motion, and that its mass was purely electromagnetic, i.e., the result of interaction with its self-field. This is Lorentz's version of the classical dream referred to by Pais in the passage we quoted in the introduction. The mass-velocity relations for Lorentz's electron model are just the relativistic relations (2.31)–(2.32). So it is indeed no coincidence that Lorentz found these same relations twice, first, in 1899, as a necessary condition for rendering ether drift unobservable (see eqs. (3.8)–(3.14)) and then, in 1904, as the mass-velocity relations for a concrete Lorentz invariant model of the electron. But the explanation is not, as Lorentz thought, that his model provides an accurate representation of the real electron; it is simply that the mass of *any* Lorentz-invariant model of *any* particle—whatever its nature and whatever its shape—will exhibit the exact same velocity dependence. This was first shown (for static systems) by Laue (1911a) and, to use Pais' imagery again, it killed Lorentz's dream.

Quite independently of Laue's later analysis, Lorentz's electron model appeared to be dead on arrival. The model as it stands is inconsistent. One way to show this is to compare expression (4.21) for the longitudinal mass $m_{//}$ derived from the electron's electromagnetic momentum to the expression for $m_{//}$ derived from its electromagnetic energy (see eq. (4.7)). These two calculations, it turns out, do not give the same result (Abraham 1905, 188, 204).⁴⁰ Setting $l = 1$ in eq. (4.18) and using eq. (2.41), we find

$$m_{//} = \frac{1}{v} \frac{4}{3} \frac{d\gamma}{dv} U'_{\text{EM}} \pm \frac{1}{3v} \frac{d}{dv} \left(\frac{1}{\gamma} \right) U'_{\text{EM}} = \gamma^3 \frac{4}{3} \frac{U'_{\text{EM}}}{c^2} \pm \frac{1}{3v} \frac{d}{dv} \left(\frac{1}{\gamma} \right) U'_{\text{EM}}. \quad (4.22)$$

The first term in the last expression is equal to $m_{//}$ in eq. (4.19) for $l = 1$. Without even working out the second term, we thus see that momentum and energy lead to different expressions for the longitudinal mass of Lorentz's electron.

For the Bucherer-Langevin electron there is no ambiguity in the formula for its longitudinal mass. Inserting $l = \gamma^{-1/3}$ and eq. (4.20) into eq. (4.19), we find that the electromagnetic momentum of the Bucherer-Langevin electron gives:

$$m_{//} = \frac{dP_{\text{EM}}}{dv} = \left(\gamma^{8/3} + \gamma v \frac{d\gamma^{-1/3}}{dv} \right) m_0 = \gamma^{8/3} \left(1 - \frac{1}{3} \beta^2 \right) m_0, \quad (4.23)$$

⁴⁰ Cf. (Miller 1981, sec. 1.13.2). Miller cites a letter of January 26, 1905, in which Abraham informed Lorentz of this difficulty. See also (Lorentz 1915, 213).

where in the last step we used eq. (2.41) in conjunction with

$$\frac{d\gamma^{-1/3}}{dv} = -\frac{1}{3}\gamma^{-4/3}\frac{d\gamma}{dv} = -\frac{1}{3}\gamma^{5/3}\frac{v}{c^2}. \quad (4.24)$$

Inserting $l = \gamma^{-1/3}$ and eq. (4.20) into eq. (4.18), we find that its electromagnetic energy gives:

$$m_{//} = \frac{1}{v} \frac{dU_{EM}}{dv} = \frac{c^2}{v} \frac{d}{d\gamma} \left(\gamma^{2/3} - \frac{1}{4}\gamma^{-4/3} \right) \frac{d\gamma}{dv} m_0. \quad (4.25)$$

Some simple gamma gymnastics establishes that eq. (4.25) reproduces eq. (4.23):⁴¹

$$m_{//} = \frac{1}{v} \frac{dU_{EM}}{dv} = \gamma^{8/3} \left(1 - \frac{1}{3}\beta^2 \right) m_0 \quad (4.26)$$

So the energy and the momentum of the Bucherer-Langevin electron do indeed give the same longitudinal mass. The same is true for the Abraham electron, although the calculation is more involved and unimportant for our purposes.

One thing the Abraham model and the Bucherer-Langevin model have in common and what distinguishes both models from Lorentz's is that the volume of the electron is constant. Hence, whatever forces are responsible for stabilizing the electron never do any work and can safely be ignored, as was done in the derivation of the basic equations (2.32) and (2.37) for longitudinal mass (see eqs. (2.13) and (2.35) and notes 21 and 26). This does not mean that no such forces are needed. In all three models, one is faced with the problem of the electron's stability. Abraham, however, argued that whereas Lorentz's contractile electron called for the explicit addition of non-electromagnetic stabilizing forces, he, Abraham, could simply take the rigidity of his own spherical electron as a given and proceed from there without ever running into trouble.

⁴¹ Eq. (4.25) can be rewritten as:

$$m_{//} = \frac{c^2}{v} \left(\frac{2}{3}\gamma^{-1/3} + \frac{1}{3}\gamma^{-7/3} \right) \gamma^3 \frac{v}{c^2} m_0.$$

Where we used eq. (2.41) for $d\gamma/dv$. This in turn can be rewritten as

$$m_{//} = \gamma^{8/3} \left(\frac{2}{3} + \frac{1}{3}\gamma^{-2} \right) m_0 = \gamma^{8/3} \left(\frac{2}{3} + \frac{1}{3}(1-\beta^2) \right) m_0 = \gamma^{8/3} \left(1 - \frac{1}{3}\beta^2 \right) m_0.$$

In the introduction of the 1903 exposition of his electron dynamics, Abraham (1903, 108–109) devoted two long paragraphs to the justification of this crucial assumption. He distinguished three sets of equations for the dynamics of the electron. We already encountered two of these, the “field equations” determining the self-field of the electron and the “fundamental dynamical equations” determining the motion of the electron in an external field. Logically prior to these, however, is what Abraham called the “basic kinematical equation,” which “limits the freedom of motion of the electron.” This is the assumption that the electron always retains its spherical shape. Abraham tried to preempt the criticism he anticipated on this score:

This basic kinematical hypothesis may strike many as arbitrary; invoking the analogy with ordinary electrically charged solid bodies, many would subscribe to the view that the truly enormous field strengths at the surface of the electron—field strengths a trillion times larger than those amenable to measurement—are capable of deforming the electron; that electrical and elastic forces on a spherical electron would be in equilibrium as long as the electron is at rest; but that the motion of the electron would change the forces of the electromagnetic field, and thereby the shape of the equilibrium state of the electron. This is not the view that has led to agreement with experiment. It also seemed to me that the assumption of a deformable electron is not allowed on fundamental grounds. The assumption leads to the conclusion that work is done either by or against the electromagnetic forces when a change of shape takes place, which means that in addition to the electromagnetic energy an internal potential energy of the electron needs to be introduced. If this were really necessary, it would immediately make an electromagnetic foundation of the theory of cathode and Becquerel rays, purely electric phenomena, impossible: one would have to give up on an electromagnetic foundation of mechanics right from the start. It is our goal, however, to provide a purely electromagnetic foundation for the dynamics of the electron. For that reason we are no more entitled to ascribe elasticity to the electron than we are to ascribe material mass to it. On the contrary, our hope is to learn to understand the elasticity of matter on the basis of the electromagnetic conception (Abraham 1903, 108–109).

The suggestion that experimental data, presumably those of Kaufmann, supported his kinematics was wishful thinking on Abraham’s part (cf. Miller 1981, secs. 1.9 and 1.11). In support of his more general argument, Abraham proceeded to appeal to no less an authority than Heinrich Hertz:

Hertz has convincingly shown that one is allowed to talk about rigid connections before one has talked about forces. Our dynamics of the electron does not talk about forces trying to deform the electron at all. It only talks about “external forces,” which try to give [the electron] a velocity or an angular velocity, and about “internal forces,” which stem from the [self-]field of the electron and which balance these external forces. Even these “forces” and “torques” are only auxiliary quantities defined in terms of the fundamental kinematic and electromagnetic concepts. The same holds for terms like “work,” “energy,” and “momentum.” The guiding principle in choosing these terms, however, was to bring out clearly the analogy between electromagnetic mechanics and the ordinary mechanics of material bodies (Abraham 1903, 109).

Abraham submitted this paper in October 1902, almost three years before the publication of Einstein's first paper on relativity. He can thus hardly be faulted for basing his new electromagnetic mechanics on the old Newtonian kinematics. Minkowski would sneer a few years later that "approaching Maxwell's equation with the concept of a rigid electron seems to me the same thing as going to a concert with your ears stopped up with cotton wool" (quoted in Miller 1973, sec. 12.4.5, 330). He made this snide comment during the 80th *Versammlung Deutscher Naturforscher und Ärzte* in Cologne in September 1908, the same conference where he gave his now famous talk "Space and Time" (Minkowski 1909). His veritable diatribe against the rigid electron, which he called a "monster" and "no working hypothesis but a working hindrance," came during the discussion following a talk by Bucherer (1908c), who presented data that seemed to contradict Abraham's predictions for the velocity dependence of electron mass and support what was by then no longer just Lorentz's prediction but Einstein's as well. It was only decades later that these data were also shown to be inconclusive (Zahn and Spees 1938; quoted in Miller 1981, 331).

Minkowski's comment suggests that we run Abraham's argument about the kinematics of the electron in Minkowski rather than in Newtonian space-time. We would then take it as a given that the electron has the shape of a sphere in its rest frame, which implies that it will have the shape of a sphere contracted in the direction of motion in any frame in which it is moving. This, of course, is exactly Lorentz's model. This gives rise to a little puzzle. The point of Abraham's argument in the passage we just quoted was that by adopting rigid kinematical constraints we can safely ignore stabilizing forces. His objection to Lorentz's model was that Lorentz *did* have to worry about non-electromagnetic stabilizing forces or he would end up with two different formula for the longitudinal mass of his electron. How can these two claims by Abraham be reconciled with one another? One's initial reaction might be that Abraham's kinematical argument does not carry over to special relativity because the theory leaves no room for rigid bodies. That in and of itself is certainly true, but it is not the source of the problem. We could run the argument using some appropriate concept of an *approximately* rigid body. Abraham's argument that kinematic constraints can be used to obviate the need for discussion of the stability of the electron will then go through *as long as we use proper relativistic notions*. From a relativistic point of view, the analysis of Lorentz's model in this section is based on the standard non-covariant definition (2.9) of four-momentum. If we follow Fermi, Rohrlich and others and use definition (2.11) (with a fixed unit vector n^μ) instead, the ambiguity in the longitudinal mass of Lorentz's electron simply disappears. After all, under this alternative definition the combination of the energy and

momentum of the electron's self-field transforms as a four-vector, even though it is an open system. This, in turn, guarantees—as we saw in eqs. (2.38)–(2.43) at the end of sec. 2—that energy and momentum give the same longitudinal mass. This shows that the ambiguity in the longitudinal mass of Lorentz's electron is not a consequence of the instability of the electron, but an artifact of the definitions of energy and momentum he used. We do not claim great originality for this insight. It is simply a matter of translating Rohrlich's analysis of the “4/3 puzzle” in special relativity (see the discussion following eq. (4.15)) to a pre-relativistic setting.

4.5 Hamiltonian, Lagrangian, and generalized momentum. Poincaré (1906, 524) brought out the inconsistency of Lorentz's model in a slightly different way. He raised the question whether the expressions he found for energy, momentum, and Lagrangian for the field of the moving electron conform to the standard relations between Hamiltonian, Lagrangian, and generalized momentum. For an electron moving in the positive x -direction, these relations are

$$U = \mathbf{P} \cdot \mathbf{v} - L = P_v - L, \quad P = P_x = \frac{dL}{dv}. \quad (4.27)$$

It turns out that the first relation is satisfied by both the Lorentz and the Bucherer-Langevin model, but that the second is satisfied only by the latter. Using eqs. (4.11) and (4.15) for P_{EM} and L_{EM} , respectively, we find

$$P_{\text{EM}}v - L_{\text{EM}} = \left(\frac{4}{3} \gamma l \beta^2 + \frac{l}{\gamma} \right) U'_{\text{EM}} = \gamma l \left(\frac{4}{3} \beta^2 + 1 - \beta^2 \right) U'_{\text{EM}}, \quad (4.28)$$

which does indeed reduce to the expression for U_{EM} found in eq. (4.6) for any value of l .

We now compute the conjugate momentum.

$$\frac{dL_{\text{EM}}}{dv} = -U'_{\text{EM}} \frac{d}{dv} \left(\frac{l}{\gamma} \right) = -U'_{\text{EM}} \left\{ \frac{1}{\gamma} \frac{dl}{dv} - \frac{l}{\gamma^2} \frac{d\gamma}{dv} \right\} = U'_{\text{EM}} \left\{ l \gamma \frac{v}{c^2} - \frac{1}{\gamma} \frac{dl}{dv} \right\} \quad (4.29)$$

where we used eq. (2.41). For the Lorentz model, with $l = 1$, this reduces to

$$\frac{dL_{\text{EM}}}{dv} = \gamma \left(\frac{U'_{\text{EM}}}{c^2} \right) v \quad (4.30)$$

which differs by the now familiar factor of $4/3$ from the expression for P_{EM} read off from eq. (4.15) for $l = 1$. For the Bucherer-Langevin model, $l = \gamma^{-1/3}$ and with the help of eq. (4.24) we find:

$$\frac{dL_{\text{EM}}}{dv} = \frac{U'_{\text{EM}}}{c^2} \left\{ \gamma^{2/3} v + \frac{1}{3} \gamma^{2/3} v \right\} = \frac{4}{3} \gamma^{2/3} \left(\frac{U'_{\text{EM}}}{c^2} \right) v. \quad (4.31)$$

This agrees exactly with eq. (4.15) for $l = \gamma^{-1/3}$.

The relations (4.27) are automatically satisfied if $(U/c, \mathbf{P})$ transforms as a four-vector under Lorentz transformations. In that case, we have (see eq. (2.6)):

$$U = \gamma U_0, \quad P = \gamma \frac{U_0}{c^2} v. \quad (4.32)$$

Inserting this into $L = Pv - U$, we find

$$L = \gamma U_0 \beta^2 - \gamma U_0 = -\gamma U_0 (1 - \beta^2) = -\frac{U_0}{\gamma}, \quad (4.33)$$

which, in turn, implies that

$$\frac{dL}{dv} = -U_0 \frac{d}{dv} \left(\frac{1}{\gamma} \right) = \frac{U_0}{\gamma^2} \frac{d\gamma}{dv} = \gamma \frac{U_0}{c^2} v, \quad (4.34)$$

in accordance with eq. (4.32). This shows once again (cf. the discussion at the end of sec. 4.4) that the inconsistency in Lorentz's model can be taken care of by switching—in relativistic terms—from the standard definition (2.9) of four-momentum to the Fermi-Rohrlich definition (2.11) (with fixed n^μ). In that case the energy and momentum of the electron's self-field will satisfy eqs. (4.32)–(4.34) even though it is an open system.

5. Poincaré pressure.

In this section we give a streamlined version of the argument with which Poincaré (1906, 525–529) introduced what came to be known as “Poincaré pressure” to stabilize Lorentz’s purely electromagnetic electron.⁴²







| | real electron (in motion) dimensions: $(r, \vartheta r, \vartheta r)$ | ideal electron (at rest) dimensions: $(\gamma lr, l\vartheta r, l\vartheta r)$ |
|---|---|--|
| The rigid electron of Abraham $\vartheta = 1$ l arbitrary r constant |  sphere (r, r, r) |  ellipsoid $(\gamma lr, lr, lr)$ |
| The contractile electron of Lorentz and Poincaré $\vartheta = \gamma$ $l = 1$ $\gamma r = \text{constant}$ |  ellipsoid $(r, \gamma r, \gamma r)$ |  sphere $(\gamma r, \gamma r, \gamma r)$ |
| The contractile electron of constant volume of Bucherer and Langevin $\vartheta = \gamma$ $l = \gamma^{-1/3}$ $\gamma lr = \text{constant}$ |  ellipsoid $(r, \gamma r, \gamma r)$ |  sphere $(\gamma^{2/3}r, \gamma^{2/3}r, \gamma^{2/3}r)$ |

Figure 2: Poincaré’s characterization of the different models for a moving electron and the corresponding states at rest in the ether.

⁴² We are grateful to Serge Rudaz for his help in reconstructing this argument.

The Lagrangian for the electromagnetic field of a moving electron can in all three models (Abraham, Lorentz, Bucherer-Langevin) be written as

$$L_{\text{EM}} = \frac{\varphi(\vartheta/\gamma)}{\gamma^2 r} \quad (5.1)$$

(Poincaré 1906, 525), where the argument ϑ/γ of the as yet unknown function φ is the ‘ellipticity’ (our term) of the “ideal electron” (Poincaré’s term for the corresponding state of the moving electron). The ellipticity is the ratio of the radius of the “ideal electron” in the directions perpendicular to the direction of motion ($l\vartheta r$) and its radius ($\gamma l r$) in the direction of motion. This is illustrated in Fig. 2, which is the same as Fig. 1, except that it shows the notation Poincaré used to describe the three electron models.

For the Abraham electron the ellipticity is $1/\gamma$; for both the Lorentz and the Bucherer-Langevin electron it is 1. By examining the Lorentz case, we can determine $\varphi(1)$. Inserting $U_{0\text{EM}} = e^2/8\pi\epsilon_0\gamma r$,⁴³ where γr is the radius of the electron at rest in the ether, into eq. (4.11) for the Lagrangian, we find:

$$L_{\text{EM Lorentz}} = -\frac{U_{0\text{EM}}}{\gamma} = -\frac{e^2}{8\pi\epsilon_0\gamma^2 r}, \quad (5.2)$$

Comparison with the general expression for L_{EM} in eq. (5.1) gives:

$$\varphi(1) = -\frac{e^2}{8\pi\epsilon_0}. \quad (5.3)$$

Abraham (1902a, 37) found that the Lagrangian for his electron model has the form

$$L_{\text{EM Abraham}} = \frac{a}{r} \frac{1-\beta^2}{\beta} \ln \frac{1+\beta}{1-\beta} \quad (5.4)$$

(Poincaré 1906, 526). Since $L_{\text{EM}} = \varphi(1)/r$ for $\beta = 0$ (in which case all three electron models coincide), it follows that $a = \varphi(1)$. From eqs. (5.1) and (5.4) it follows that

$$\varphi(1/\gamma) = \gamma^2 r L_{\text{EM Abraham}} = \frac{a}{\beta} \ln \frac{1+\beta}{1-\beta}. \quad (5.5)$$

⁴³ See note 48 below for a derivation of this formula.

The Lagrangian for the Lorentz model told us that $\varphi(1) = a$. The Lagrangian for the Abraham model allows us to determine $\varphi'(1)$. We start from eq. (5.5) and develop both the right-hand side and the argument $1/\gamma$ of φ on the left-hand side to second order in β . This gives (ibid.):

$$\varphi\left(1 - \frac{1}{2}\beta^2\right) = a\left(1 + \frac{1}{3}\beta^2\right). \quad (5.6)$$

Now differentiate both sides:

$$-\beta\varphi'\left(1 - \frac{1}{2}\beta^2\right) = \frac{2}{3}a\beta. \quad (5.7)$$

It follows that $\varphi'(1) = -(2/3)a$.

As Poincaré notes, all three electron models satisfy a constraint of the form

$$r = b\vartheta^m \quad (5.8)$$

where b is a constant and where the exponent m depends on which model we consider. In the Abraham model $\vartheta = 1$ and r is a constant. Hence, $r = b$. In the Lorentz model, $\vartheta = \gamma$ and γr is a constant. It follows that $\vartheta r = b$, or $r = b\vartheta^{-1}$. In the Bucherer-Langevin model, $\vartheta = \gamma$ and $\gamma^{2/3}r$ is a constant. It follows that $\vartheta^{2/3}r = b$, or $r = b\vartheta^{-2/3}$. In other words, the values of m in the three models are

$$\text{Abraham: } m = 0$$

$$\text{Lorentz: } m = -1 \quad (5.9)$$

$$\text{Bucherer-Langevin: } m = -2/3$$

Substituting $r = b\vartheta^m$ into the general expression (5.1) for the Lagrangian, we find:

$$L_{\text{EM}} = \frac{\varphi(\vartheta/\gamma)}{b\gamma^2\vartheta^m}. \quad (5.10)$$

Poincaré proceeds to investigate whether this Lagrangian describes a stable physical system. To this end, he checks whether $\partial L_{\text{EM}}/\partial\vartheta$ vanishes. It turns out that for $m = -2/3$ it does, but that for $m = -1$ it does not. Denote the argument of the function φ with $u \equiv \vartheta/\gamma$.

$$\frac{\partial L_{\text{EM}}}{\partial\vartheta} = \frac{\varphi'(u)}{b\gamma^3\vartheta^m} - \frac{m\varphi(u)}{b\gamma^2\vartheta^{m+1}}. \quad (5.11)$$

This derivative vanishes if

$$\varphi'(u) = \frac{\gamma m \varphi(u)}{\vartheta} = m \frac{\varphi(u)}{u} \quad (5.12)$$

For the Lorentz and Bucherer-Langevin models $u = 1$, and this condition reduces to

$$\varphi'(1) = m\varphi(1). \quad (5.13)$$

Inserting $\varphi(1) = a$ and $\varphi'(1) = -(2/3)a$, we see that the purely electromagnetic Lagrangian only describes a stable system for $m = -2/3$, which is the value for the Bucherer-Langevin electron. The Lorentz electron calls for an additional term in the Lagrangian. The total Lagrangian is then given by the sum

$$L_{\text{tot}} = L_{\text{EM}} + L_{\text{non-EM}}. \quad (5.14)$$

Like L_{EM} , $L_{\text{non-EM}}$ is a function of ϑ and r . Treating these variables as independent, we can write the stability conditions for the total Lagrangian as

$$\frac{\partial}{\partial \vartheta} (L_{\text{EM}} + L_{\text{non-EM}}) = 0, \quad \frac{\partial}{\partial r} (L_{\text{EM}} + L_{\text{non-EM}}) = 0. \quad (5.15)$$

Evaluating the partial derivatives of L_{EM} given by eq. (5.1),

$$\frac{\partial L_{\text{EM}}}{\partial \vartheta} = \frac{\varphi'(u)}{\gamma^3 r}, \quad \frac{\partial L_{\text{EM}}}{\partial r} = -\frac{\varphi(u)}{\gamma^2 r^2}, \quad (5.16)$$

and inserting the results into the stability conditions, we find

$$\frac{\partial L_{\text{non-EM}}}{\partial \vartheta} = -\frac{\varphi'(u)}{\gamma^3 r}, \quad \frac{\partial L_{\text{non-EM}}}{\partial r} = \frac{\varphi(u)}{\gamma^2 r^2}. \quad (5.17)$$

Poincaré (1906, 528–529) continues his analysis without picking a specific model. We shall only do the calculation for the Lorentz model. So we no longer need subscripts such as in eqs. (5.2) and (5.4) to distinguish between the models of Abraham and Lorentz. For the Lorentz model $m = -1$, $\gamma = \vartheta$, $r = b/\vartheta$, and $u = 1$. Substituting these values into eqs. (5.17) and using that $\varphi(1) = a$ and $\varphi'(1) = -(2/3)a$, we find:

$$\frac{\partial L_{\text{non-EM}}}{\partial \vartheta} = \frac{2a}{3b\vartheta^2}, \quad \frac{\partial L_{\text{non-EM}}}{\partial r} = \frac{a}{b^2}. \quad (5.18)$$

These equations are satisfied by a Lagrangian of the form

$$L_{\text{non-EM}} = Ar^3\vartheta^2, \quad (5.19)$$

where A is a constant. Since $r^3\vartheta^2$ is proportional to the volume V of the moving electron, $L_{\text{non-EM}}$ can be written as

$$L_{\text{non-EM}} = P_{\text{Poincaré}}V \quad (5.20)$$

where $P_{\text{Poincaré}}$ is a constant. We chose the letter P because this constant turns out to be a (negative) pressure. To determine the constant A , we take the derivative of eq. (5.19) with respect to ϑ and r , and eliminate r from the results, using $r = b/\vartheta$:

$$\frac{\partial L_{\text{non-EM}}}{\partial \vartheta} = 2Ar^3\vartheta = \frac{2Ab^3}{\vartheta^2}, \quad \frac{\partial L_{\text{non-EM}}}{\partial r} = 3Ar^2\vartheta^2 = 3Ab^2. \quad (5.21)$$

Comparison with eqs. (5.18) gives:

$$A = \frac{a}{3b^4}. \quad (5.22)$$

Finally, we write $L_{\text{non-EM}}$ in a form that allows easy comparison with $L_{\text{EM}} = a/\gamma^2 r$ (see eq. (5.1) with $\varphi(\vartheta/\gamma) = \varphi(1) = a$). Using eq. (5.22) along with $\vartheta = \gamma$ and $b = \gamma r$, we can rewrite eq. (5.19) for $L_{\text{non-EM}}$ as

$$L_{\text{non-EM}} = \frac{a}{3b^4} r^3 \vartheta^2 = \frac{1}{3} \frac{a}{\gamma^2 r} = \frac{1}{3} L_{\text{EM}} = -\frac{1}{3} \frac{U_{0\text{EM}}}{\gamma}, \quad (5.23)$$

where in the last step we used eq. (4.11) for $l = 1$. Using that the volume V_0 of Lorentz's electron at rest is equal to γV , we can rewrite this as:

$$L_{\text{non-EM}} = -\frac{1}{3} \frac{U_{0\text{EM}}}{V_0} V, \quad (5.24)$$

Comparison with expression (5.20) gives:

$$P_{\text{Poincaré}} = -\frac{1}{3} \frac{U_{0\text{EM}}}{V_0} \quad (5.25)$$

(Laue 1911b, 164, eq. 171). Note that this so-called Poincaré pressure is negative. Although Poincaré does not explicitly state this, it is clear from the context that the

pressure is present only inside the electron and vanishes outside. It can be written more explicitly with the help of the ϑ -step-function:⁴⁴

$$P_{\text{Poincaré}}(\mathbf{x}) = -\frac{1}{3} \frac{U_{0\text{EM}}}{V_0} \vartheta\left(R - \sqrt{\gamma^2 x^2 + y^2 + z^2}\right), \quad (5.26)$$

where R is the radius of the electron at rest. So there is a sudden drop in pressure at the edge of the electron, which is the only place where forces are exerted.⁴⁵ These forces serve two purposes. First, they prevent the electron's surface charge distribution from flying apart under the influence of the Coulomb repulsion between its parts. Second, as the region where $P_{\text{Poincaré}}(\mathbf{x})$ is non-vanishing always coincides with the ellipsoid-shaped region occupied by the moving electron, these forces make the electron contract by a factor γ in the direction of motion when it is moving with a velocity v .

Relations (4.27) between Hamiltonian, Lagrangian and generalized momentum, only one of which was satisfied by Lorentz's original purely electromagnetic electron model, are both satisfied once $L_{\text{non-EM}}$ is added to the Lagrangian. Using the total Lagrangian,

$$L_{\text{tot}} = L_{\text{EM}} + L_{\text{non-EM}} = \frac{4}{3} L_{\text{EM}} = -\frac{4}{3} \frac{U_{0\text{EM}}}{\gamma}, \quad (5.27)$$

to compute the total momentum, we find:

$$P_{\text{tot}} = \frac{dL_{\text{tot}}}{dv} = \frac{4}{3} \frac{dL_{\text{EM}}}{dv} = \frac{4}{3} \gamma \frac{U_{0\text{EM}}}{c^2} v, \quad (5.28)$$

where in the last step we used eq. (4.30). This is just the electromagnetic momentum P_{EM} found earlier (see eq. (4.15) for $l=1$). With the help of these expressions for L_{tot} and P_{tot} , we can compute the total energy.

$$U_{\text{tot}} = P_{\text{tot}} v - L_{\text{tot}} = \frac{4}{3} \gamma U_{0\text{EM}} \beta^2 + \frac{4}{3} \frac{U_{0\text{EM}}}{\gamma} = \frac{4}{3} \gamma U_{0\text{EM}}. \quad (5.29)$$

The total energy is the sum of the electromagnetic energy (see eq. (4.7)),

⁴⁴ The definition of this function is: $\vartheta(x)=0$ for $x<0$ and $\vartheta(x)=1$ for $x\geq 0$.

⁴⁵ For a detailed analysis of the completely analogous case of the forces on a capacitor in the Trouton-Noble experiment, see secs. 2.3.3 and 2.4.2 of (Janssen 1995).

$$U_{\text{EM}} = \frac{4}{3}\gamma U_{0\text{EM}} - \frac{1}{3}\frac{U_{0\text{EM}}}{\gamma}, \quad (5.30)$$

and the non-electromagnetic energy,

$$U_{\text{non-EM}} = \frac{1}{3}\frac{U_{0\text{EM}}}{\gamma}, \quad (5.31)$$

which is minus the product of the Poincaré pressure (see eq. (5.25)) and the volume $V = V_0/\gamma$ of the moving electron. The total energy of the system at rest is

$$U_{0\text{tot}} = \frac{4}{3}U_{0\text{EM}}. \quad (5.32)$$

and its rest mass is $m_{0\text{tot}} = U_{0\text{tot}}/c^2$ accordingly. Eq. (5.28) can thus be rewritten as

$$P_{\text{tot}} = \gamma \left(\frac{U_{0\text{tot}}}{c^2} \right) v = \gamma m_{0\text{tot}} v. \quad (5.33)$$

The troublesome factor 4/3 has disappeared.

The total energy and momentum transform as a four-vector under Lorentz transformations. In the system's rest frame its four-momentum is $P_{0\text{tot}}^\mu = (U_{0\text{tot}}/c, 0, 0, 0)$. In a frame moving with velocity v in the x -direction, it is

$$P_{\text{tot}}^\mu = \Lambda^\mu{}_\nu P_{0\text{tot}}^\nu = \left(\gamma \frac{U_{0\text{tot}}}{c}, \gamma \beta \frac{U_{0\text{tot}}}{c}, 0, 0 \right) \quad (5.34)$$

in accordance with eqs. (5.29), (5.32), and (5.33). As we saw at the end of sec. 2, if $(U/c, \mathbf{P})$ transforms as a four-vector, it is guaranteed that energy and momentum lead to the same longitudinal mass. With Poincaré's amendment Lorentz's electron model may no longer be purely electromagnetic—at least it is fully consistent.

6. The relativistic treatment of the electron model of Lorentz as amended by Poincaré.

From the point of view of Laue's relativistic continuum mechanics, the problem with Lorentz's fully electromagnetic electron is that it is not a closed system. The four-divergence of the energy-momentum tensor of its electromagnetic field does not vanish. Computing this four-divergence tells us what needs to be added to this energy-momentum tensor to obtain a closed system, i.e., a system with a total energy-momentum tensor such that $\partial_\nu T_{\text{tot}}^{\mu\nu} = 0$. Unsurprisingly, the part that needs to be added is just the energy-momentum tensor for the Poincaré pressure.

The energy-momentum tensor for the electromagnetic field is given by

$$T_{\text{EM}}^{\mu\nu} = \mu_0^{-1} \left(F^\mu{}_\alpha F^{\alpha\nu} + \frac{1}{4} \eta^{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} \right), \quad (6.1)$$

where $F^{\mu\nu}$ is the electromagnetic field tensor with components:

$$F^{\mu\nu} = \begin{pmatrix} 0 & -E_x/c & -E_y/c & -E_z/c \\ E_x/c & 0 & -B_z & B_y \\ E_y/c & B_z & 0 & -B_x \\ E_z/c & -B_y & B_x & 0 \end{pmatrix}. \quad (6.2)$$

Compared to $F^{\mu\nu}$ the $0i$ and ij components of $F^\mu{}_\nu = F^{\mu\alpha} \eta_{\alpha\nu}$ have the opposite sign, as do the $0i$ and $i0$ components of $F_{\mu\nu} = \eta_{\mu\alpha} F^{\alpha\beta} \eta_{\beta\nu}$. Inserting the components of the field tensor into eq. (6.1) for the energy-momentum tensor, we recover the familiar expressions for the electromagnetic energy density (cf. eq. (4.2)), (c times) the electromagnetic momentum density (cf. eq. (2.15)), and (minus) the Maxwell stress tensor (cf. eq. (2.25)).

$$\begin{aligned} T_{\text{EM}}^{00} &= \frac{1}{2} \varepsilon_0 E^2 + \frac{1}{2} \mu_0^{-1} B^2 = u_{\text{EM}}, \\ \left(T_{\text{EM}}^{01}, T_{\text{EM}}^{02}, T_{\text{EM}}^{03} \right) &= \left(T_{\text{EM}}^{10}, T_{\text{EM}}^{20}, T_{\text{EM}}^{30} \right) = c \varepsilon_0 \mathbf{E} \times \mathbf{B} = c \mathbf{p}_{\text{EM}}, \\ T_{\text{EM}}^{ij} &= -\varepsilon_0 \left(E^i E^j - \frac{1}{2} \delta^{ij} E^2 \right) - \mu_0^{-1} \left(B^i B^j - \frac{1}{2} \delta^{ij} B^2 \right) = -T_{\text{Maxwell}}^{ij}. \end{aligned} \quad (6.3)$$

We calculate the four-divergence of the energy-momentum tensor for the electromagnetic field of Lorentz's electron in its rest frame. Lorentz invariance

guarantees that if the four-divergence of the total energy-momentum tensor vanishes in the rest frame ($\partial_{0\nu} T_{\text{tot}}^{\mu\nu} = 0$), it will vanish in all frames ($\partial_\nu T_{\text{tot}}^{\mu\nu} = 0$). In the rest frame, we have

$$T_{0\text{EM}}^{\mu\nu} = \begin{pmatrix} u_{0\text{EM}} & 0 \\ 0 & -T_{0\text{Maxwell}}^{ij} \end{pmatrix}. \quad (6.4)$$

Consider the four-divergence $\partial_{0\nu} T_{0\text{EM}}^{\mu\nu}$ of this tensor. Since the system is static, only the spatial derivatives, $\partial_{0j} T_{0\text{EM}}^{ij}$, give a contribution. Since $T_{0\text{EM}}^{0j} = 0$, there will only be contributions for $\mu = i$. Going through eqs. (2.20)–(2.25) in reverse order and setting $\mathbf{B} = 0$, we can write these contributions as:

$$\partial_{0j} T_{0\text{EM}}^{ij} = -\partial_{0j} T_{0\text{Maxwell}}^{ij} = -\varepsilon_0 E_0^i \text{div} \mathbf{E}_0 = -\rho_0 E_0^i, \quad (6.5)$$

where in the last step we used one of Maxwell's equations.⁴⁶ The charge density ρ_0 is the surface charge density $\sigma = e/4\pi R^2$ (where e is the charge of the electron and R the radius of the electron in its rest frame):

$$\rho_0 = \sigma \delta(R - r_0), \quad (6.6)$$

where $r_0 \equiv \sqrt{x_0^2 + y_0^2 + z_0^2}$. Inside the electron there is no electric field (it is a miniature version of Faraday's cage); outside the field is the same as that of a point charge e located at the center of the electron. We thus have:

$$E_0^i = \begin{cases} r_0 < R: & 0 \\ r_0 > R: & \frac{e}{4\pi\varepsilon_0 r_0^2} \frac{x_0^i}{r_0}. \end{cases} \quad (6.7)$$

At $r_0 = R$, right on the surface of the electron, the field has a discontinuity. Its magnitude, E_0 , jumps from 0 to $e/4\pi\varepsilon_0 R^2$. At this point we need to use the average of these two values (see, e.g., Griffith 1999, 102–103). At $r_0 = R$ the field is thus given by

⁴⁶ From a relativistic point of view, eq. (6.5) is immediately obvious since the (four-)gradient of the energy-momentum tensor gives minus the density of the (four-)force acting on the system (see, e.g., Pauli 1921, 126, eq. (345)). The right-hand side of eq. (6.5) is minus the Lorentz force density in the absence of a magnetic field (ibid., 85, eq. (225)).

$$E_{r_0=R}^i = \frac{e}{8\pi\epsilon_0 R^2} \frac{x_0^i}{R} = \frac{\sigma}{2\epsilon_0} \frac{x_0^i}{R}, \quad (6.8)$$

where we used that $\sigma = e/4\pi R^2$. Substituting eqs. (6.8) and (6.6) into eq. (6.5), we find:

$$\partial_{0_j} T_{0_{EM}}^{ij} = -\frac{\sigma^2}{2\epsilon_0} \frac{x_0^i}{R} \delta(R - r_0). \quad (6.9)$$

In summary, the divergence of the energy-momentum tensor of the electron's electromagnetic field in its rest frame is:

$$\partial_{0_\nu} T_{0_{EM}}^{\mu\nu} = \begin{cases} \mu = 0: & 0 \\ \mu = i: & -\frac{\sigma^2}{2\epsilon_0} \frac{x_0^i}{R} \delta(R - r_0). \end{cases} \quad (6.10)$$

It vanishes everywhere except at the surface of the electron where its charge is. To get a total energy-momentum tensor with a four-divergence that vanishes everywhere,

$$\partial_{0_\nu} T_{0_{tot}}^{\mu\nu} = \partial_{0_\nu} \left(T_{0_{EM}}^{\mu\nu} + T_{0_{non-EM}}^{\mu\nu} \right) = 0, \quad (6.11)$$

we need to add the Poincaré pressure of eq. (5.26), which in the electron's rest frame is described by the energy-momentum tensor

$$T_{0_{non-EM}}^{\mu\nu} = -\eta^{\mu\nu} P_{\text{Poincaré}} \vartheta(R - r_0). \quad (6.12)$$

We calculate the four-divergence of this energy-momentum tensor, $\partial_{0_\nu} T_{0_{non-EM}}^{\mu\nu}$. Only the ij -components will contribute (cf. eqs. (6.4)–(6.5) above):⁴⁷

$$\partial_{0_j} T_{0_{non-EM}}^{ij} = P_{\text{Poincaré}} \partial_{0_i} \vartheta(R - r_0) = -P_{\text{Poincaré}} \frac{x_0^i}{R} \delta(R - r_0). \quad (6.13)$$

The divergence of the energy-momentum tensor describing the Poincaré pressure in the electron's rest frame is thus given by:

⁴⁷ This shows that we have considerable freedom in choosing the 00-component of $T_{0_{non-EM}}^{\mu\nu}$. Instead of $-\vartheta(R - r_0) P_{\text{Poincaré}}$ we can pick *any* function of the spatial coordinates. The most convenient choice is $T_{0_{non-EM}}^{00} = 0$.

$$\partial_{0\nu} T_0^{\mu\nu}{}_{\text{non-EM}} = \begin{cases} \mu = 0: & 0 \\ \mu = i: & -P_{\text{Poincaré}} \frac{x_0^i}{R} \delta(R - r_0). \end{cases} \quad (6.14)$$

Comparison of eqs. (6.10) and (6.14) shows that the four-divergence of the electron's total energy-momentum tensor vanishes, if

$$P_{\text{Poincaré}} = -\frac{\sigma^2}{2\varepsilon_0}. \quad (6.15)$$

This is just the value for the Poincaré pressure that we found in sec. 5 (see eq. (5.25) and Laue 1911b, 164):

$$P_{\text{Poincaré}} = -\frac{U_{0\text{EM}}}{3V_0}. \quad (6.16)$$

The rest energy $U_{0\text{EM}}$ of a sphere of radius R with surface charge e is given by:⁴⁸

$$U_{0\text{EM}} = \frac{e^2}{8\pi\varepsilon_0 R}. \quad (6.17)$$

Inserting this value into eq. (6.16), using that $V_0 = \frac{4}{3}\pi R^3$ and that $\sigma = e/4\pi R^2$, we find:

$$P_{\text{Poincaré}} = -\frac{U_{0\text{EM}}}{3V_0} = \frac{e^2}{(8\pi\varepsilon_0 R)(4\pi R^3)} = -\frac{1}{2\varepsilon_0} \left(\frac{e}{4\pi R^2} \right)^2 = -\frac{\sigma^2}{2\varepsilon_0}, \quad (6.18)$$

⁴⁸ To derive eq. (6.17), we start from:

$$U_{0\text{EM}} = \int \frac{1}{2} \varepsilon_0 E^2 d^3x = -\int \frac{1}{2} \varepsilon_0 \mathbf{E} \cdot \text{grad} \varphi d^3x.$$

Using the relation $\mathbf{E} \cdot \text{grad} \varphi = \text{div}(\mathbf{E}\varphi) - \varphi \text{div} \mathbf{E}$ along with Gauss' theorem (which tells us that the integral over $\text{div}(\mathbf{E}\varphi)$ vanishes), we can rewrite this as

$$U_{0\text{EM}} = \frac{1}{2} \varepsilon_0 \int \varphi \text{div} \mathbf{E} d^3x = \frac{1}{2} \int \varphi \rho d^3x.$$

The potential φ is equal to $e/4\pi\varepsilon_0 r$ outside the electron and to $e/4\pi\varepsilon_0 R$ inside. Using eq. (6.6) for ρ and switching to spherical coordinates, we arrive at eq. (6.17):

$$U_{0\text{EM}} = \frac{1}{2} \int \varphi \sigma \delta(R-r) r^2 \sin \vartheta dr d\vartheta d\varphi = 2\pi \left(\frac{e}{4\pi\varepsilon_0 R} \right) \left(\frac{e}{4\pi R^2} \right) R^2 = \frac{e^2}{8\pi\varepsilon_0 R}.$$

which is indeed the value we found in eq. (6.15) (see also Lorentz 1915, 214).

We now calculate the contributions of $T_{\text{EM}}^{\mu\nu}$ and $T_{\text{non-EM}}^{\mu\nu}$ to the electron's four-momentum. We begin with the contribution coming from the electron's electromagnetic field:

$$P_{\text{EM}}^{\mu} = \frac{1}{c} \int T_{\text{EM}}^{\mu 0} d^3x. \quad (6.19)$$

Using that $T^{\mu\nu} = \Lambda^{\mu}{}_{\rho} \Lambda^{\nu}{}_{\sigma} T_0^{\rho\sigma}$ and that $d^3x = d^3x_0 / \gamma$, we can rewrite this as

$$P_{\text{EM}}^{\mu} = \frac{1}{c\gamma} \Lambda^{\mu}{}_{\rho} \Lambda^0{}_{\sigma} \int T_{0\text{EM}}^{\rho\sigma} d^3x_0. \quad (6.20)$$

Eq. (6.4) tells us that there will only be contributions for $\rho\sigma = 00$ and $\rho\sigma = ij$. We denote these contributions as $P_{\text{EM}}^{\mu}(00)$ and $P_{\text{EM}}^{\mu}(ij)$.

For $P_{\text{EM}}^{\mu}(00)$ we have:

$$P_{\text{EM}}^{\mu}(00) = \frac{1}{c\gamma} \Lambda^{\mu}{}_{0} \Lambda^0{}_{0} \int T_{0\text{EM}}^{00} d^3x_0. \quad (6.21)$$

Since $\Lambda^{\mu}{}_{0} = (\gamma, \gamma\beta, 0, 0)$ (see eq. (2.4)) and the integral over $T_{0\text{EM}}^{00}$ gives $U_{0\text{EM}}$, this turns into:

$$P_{\text{EM}}^{\mu}(00) = \left(\gamma \frac{U_{0\text{EM}}}{c}, \gamma \frac{U_{0\text{EM}}}{c^2} \mathbf{v} \right). \quad (6.22)$$

This is just the Lorentz transform of $P_{0\text{EM}}^{\mu} = (U_{0\text{EM}} / c, 0, 0, 0)$. It is the additional contribution $P_{\text{EM}}^{\mu}(ij)$, coming from $T_{0\text{EM}}^{ij}$, that is responsible for the fact that the four-momentum of the electron's electromagnetic field does not transform as a four-vector.

For $P_{\text{EM}}^{\mu}(ij)$ we have:

$$P_{\text{EM}}^{\mu}(ij) = \frac{1}{c\gamma} \Lambda^{\mu}{}_{i} \Lambda^0{}_{j} \int T_{0\text{EM}}^{ij} d^3x_0. \quad (6.23)$$

The integrand is minus the Maxwell stress tensor in the electron's rest frame (see eq. (6.3)):

$$T_{0\text{EM}}^{ij} = - \begin{pmatrix} \varepsilon_0 \left(E_x^2 - \frac{1}{2} E^2 \right) & \varepsilon_0 E_x E_y & \varepsilon_0 E_x E_z \\ \varepsilon_0 E_y E_x & \varepsilon_0 \left(E_y^2 - \frac{1}{2} E^2 \right) & \varepsilon_0 E_y E_z \\ \varepsilon_0 E_z E_x & \varepsilon_0 E_z E_y & \varepsilon_0 \left(E_z^2 - \frac{1}{2} E^2 \right) \end{pmatrix}. \quad (6.24)$$

The integrals over the off-diagonal terms are all zero. The integrals over the three diagonal terms are equal to one another and given by

$$\int \varepsilon_0 \left(\frac{1}{2} E^2 - \frac{1}{3} E^2 \right) d^3 x_0 = \frac{1}{3} \int \frac{1}{2} \varepsilon_0 E^2 d^3 x_0 = \frac{1}{3} U_{0\text{EM}}. \quad (6.25)$$

With the help of this equation and of $(1/\gamma)\Lambda^\mu_i \Lambda^0_i = (1/\gamma)\Lambda^\mu_1 \Lambda^0_1 = (\gamma\beta^2, \gamma\beta, 0, 0)$ (see eq. (2.4)), eq. (6.23) can be rewritten as

$$P_{\text{EM}}^\mu(ij) = \left(\frac{1}{3} \gamma \beta^2 \frac{U_{0\text{EM}}}{c}, \frac{1}{3} \gamma \frac{U_{0\text{EM}}}{c^2} \mathbf{v} \right). \quad (6.26)$$

Adding eqs. (6.22) and (6.26), we find:

$$P_{\text{EM}}^\mu = P_{\text{EM}}^\mu(00) + P_{\text{EM}}^\mu(ij) = \left(\gamma \left(1 + \frac{1}{3} \beta^2 \right) \frac{U_{0\text{EM}}}{c}, \frac{4}{3} \gamma \frac{U_{0\text{EM}}}{c^2} \mathbf{v} \right). \quad (6.27)$$

This is exactly the result we found earlier for the energy and momentum of the electromagnetic field of Lorentz's electron (see eqs. (4.6) and (4.15) with $l=1$ and $U'_{\text{EM}} = U_{0\text{EM}}$).

The calculation of the contributions to the four-momentum coming from $T_{\text{non-EM}}^{\mu\nu}$ is completely analogous to the calculation in eqs. (6.19)–(6.27). We start with:

$$P_{\text{non-EM}}^\mu = \frac{1}{c\gamma} \Lambda^\mu_\rho \Lambda^0_\sigma \int T_{\text{non-EM}}^{\rho\sigma} d^3 x_0. \quad (6.28)$$

Since $T_{\text{non-EM}}^{\mu\nu}$ is diagonal (see eq. (6.12)), there will only be contributions when $\rho = \sigma$. Since $\Lambda^0_\mu = (\gamma, \gamma\beta, 0, 0)$, the only contributions will be for $\rho = \sigma = 0$ and $\rho = \sigma = 1$. We denote these by $P_{\text{non-EM}}^\mu(00)$ and $P_{\text{non-EM}}^\mu(11)$, respectively, and calculate them separately.

First, we consider $P_{\text{non-EM}}^\mu(00)$:

$$P_{\text{non-EM}}^\mu(00) = \frac{1}{c\gamma} \Lambda^\mu_0 \Lambda^0_0 \int T_{\text{non-EM}}^{00} d^3 x_0. \quad (6.29)$$

We can write the integral as

$$\int T_{\text{non-EM}}^{00} d^3 x_0 = -P_{\text{Poincaré}} \int \vartheta(R - r_0) d^3 x_0. \quad (6.30)$$

With eq. (6.16) for $P_{\text{Poincaré}}$ this turns into:

$$\int T_{\text{non-EM}}^{00} d^3 x_0 = \frac{U_{0\text{EM}}}{3V_0} \int \vartheta(R - r_0) d^3 x_0 = \frac{1}{3} U_{0\text{EM}}. \quad (6.31)$$

Inserting this value into eq. (6.29) along with $\Lambda^\mu_0 = (\gamma, \gamma\beta, 0, 0)$ (see eq. (2.4)), we find

$$P_{\text{non-EM}}^\mu(00) = \left(\gamma \frac{1}{3} \frac{U_{0\text{EM}}}{c}, \gamma \frac{1}{3} \frac{U_{0\text{EM}}}{c^2} \mathbf{v} \right). \quad (6.32)$$

This is the Lorentz transform of $P_{0\text{EM}}^\mu = \left(\frac{1}{3} U_{0\text{EM}} / c, 0, 0, 0 \right)$.

We now turn to $P_{\text{non-EM}}^\mu(11)$:

$$P_{\text{non-EM}}^\mu(11) = \frac{1}{c\gamma} \Lambda^\mu_1 \Lambda^0_1 \int T_{\text{non-EM}}^{11} d^3 x_0. \quad (6.33)$$

The integral over $T_{\text{non-EM}}^{11}$ can be computed in the same way as the integral over $T_{\text{non-EM}}^{00}$ in eqs. (6.30)–(6.31):

$$\int T_{\text{non-EM}}^{11} d^3 x_0 = -\frac{U_{0\text{EM}}}{3V_0} \int \vartheta(R - r_0) d^3 x_0 = -\frac{1}{3} U_{0\text{EM}}. \quad (6.34)$$

Inserting this value into eq. (6.33) along with $(1/\gamma)\Lambda^\mu_1 \Lambda^0_1 = (\gamma\beta^2, \gamma\beta, 0, 0)$, we find

$$P_{\text{non-EM}}^\mu(11) = \left(-\frac{1}{3} \gamma\beta^2 \frac{U_{0\text{EM}}}{c}, -\frac{1}{3} \gamma \frac{U_{0\text{EM}}}{c^2} \mathbf{v} \right). \quad (6.35)$$

Comparing eq. (6.35) to eq. (6.26), we see that $P_{\text{non-EM}}^\mu(11)$ is exactly the opposite of $P_{\text{EM}}^\mu(ij)$:

$$P_{\text{EM}}^\mu(ij) + P_{\text{non-EM}}^\mu(11) = 0. \quad (6.36)$$

This is a direct consequence of what is known as *Laue's theorem* (Miller 1981, 352). This theorem (Laue 1911a, 539) says that for a “complete [i.e., closed] static system” (*vollständiges statisches System*):

$$\int T_{0\text{tot}}^{ij} d^3x_0 = 0. \quad (6.37)$$

For the electron we have $T_{0\text{tot}}^{ij} = T_{0\text{EM}}^{ij} + T_{0\text{non-EM}}^{ij}$. From eqs. (6.24)–(6.25) we read off that

$$\int T_{0\text{EM}}^{ij} d^3x_0 = \begin{cases} i \neq j: & 0 \\ i = j: & \frac{1}{3}U_{0\text{EM}}. \end{cases} \quad (6.38)$$

Eq. (6.34) and similar equations for the 22- and 33-components tell us

$$\int T_{0\text{non-EM}}^{ij} d^3x_0 = \begin{cases} i \neq j: & 0 \\ i = j: & -\frac{1}{3}U_{0\text{EM}}. \end{cases} \quad (6.39)$$

Laue's theorem thus holds for this system, as it should, and eq. (6.36) is a direct consequence of this. Since $P_{\text{non-EM}}^\mu(ij) = 0$ except when $i = j = 1$, we can substitute $P_{\text{non-EM}}^\mu(ij)$ for $P_{\text{non-EM}}^\mu(11)$ in eq. (6.36). Adding this to $P_{\text{EM}}^\mu(ij)$ and using eqs. (6.20) and (6.28), we find

$$P_{\text{EM}}^\mu(ij) + P_{\text{non-EM}}^\mu(ij) = \frac{1}{c\gamma} \Lambda^\mu_i \Lambda^0_j \int \left(T_{0\text{EM}}^{ij} + T_{0\text{non-EM}}^{ij} \right) d^3x_0 \quad (6.40)$$

which by Laue's theorem vanishes, as is confirmed explicitly by eqs. (6.38)–(6.39).

Laue's theorem ensures that the four-momentum of a closed static system transforms as a four-vector. The total four-momentum of the electron is the sum of four terms (see eqs. (6.22), (6.26), (6.32), and (6.35)):

$$P_{\text{tot}}^\mu = P_{\text{EM}}^\mu(00) + P_{\text{non-EM}}^\mu(00) + P_{\text{EM}}^\mu(ij) + P_{\text{non-EM}}^\mu(11). \quad (6.41)$$

The last two terms cancel each other because of Laue's theorem, and all that is left is:

$$P_{\text{tot}}^\mu = P_{\text{EM}}^\mu(00) + P_{\text{non-EM}}^\mu(00). \quad (6.42)$$

Using eqs. (6.22) and (6.32) for these two contributions we recover eq. (5.34) for the total energy and momentum of the electron:

$$P_{\text{tot}}^\mu = \left(\gamma \frac{4}{3} \frac{U_{0\text{EM}}}{c}, \gamma \frac{4}{3} \frac{U_{0\text{EM}}}{c^2} \mathbf{v} \right) = \left(\gamma \frac{U_{0\text{tot}}}{c}, \gamma \frac{U_{0\text{tot}}}{c^2} \mathbf{v} \right). \quad (6.43)$$

As we pointed out above (see eqs. (6.11)–(6.12) and note 47), we still have a closed system if we set the 00-component of $T_{0\text{non-EM}}^{\mu\nu}$ to zero. This does not affect the result for $P_{\text{non-EM}}^\mu(11)$, which only depends on $T_{0\text{non-EM}}^{11}$ (see eq. (6.33)). $P_{\text{non-EM}}^\mu(00)$, however, will be zero if $T_{0\text{non-EM}}^{00} = 0$ (see eq. (6.29)). The total four-momentum will still be a four-vector but compared to eq. (6.43) the system's rest energy will be smaller by $\frac{1}{3}U_{0\text{EM}}$:

$$P_{\text{tot}}^\mu = P_{\text{EM}}^\mu(00) = \left(\gamma \frac{U_{0\text{EM}}}{c}, \gamma \frac{U_{0\text{EM}}}{c^2} \mathbf{v} \right). \quad (6.44)$$

To reiterate: if the stabilizing mechanism for the electron does *not* contribute to the energy in the rest frame but only to the stresses, $T_{0\text{non-EM}}^{00} = 0$ and only the first term in eq. (6.42) contributes to the four-momentum. In this case, the electron's rest mass is $m_0 = U_{0\text{EM}} / c^2$ (see eq. (6.44)). If the stabilizing mechanism *does* contribute to the energy in the electron's rest frame, $T_{0\text{non-EM}}^{00} \neq 0$ and both terms in eq. (6.42) contribute to the four-momentum. If $T_{0\text{non-EM}}^{00} = -\frac{1}{3}(U_{0\text{EM}} / V_0) \vartheta(R - r_0)$, as in Poincaré's specific model (see eqs. (6.12) and (6.16)), the electron's rest mass is $m_0 = \frac{4}{3}U_{0\text{EM}} / c^2$ (see eq. (6.43)).⁴⁹

The arbitrariness of the Lorentz-Poincaré electron is much greater than the freedom we have in choosing the 00-component of the energy-momentum tensor for the mechanism stabilizing a spherical surface charge distribution. For starters, we can choose a (surface or volume) charge distribution of any shape we like—a box, a doughnut, a banana, etc. As long as this charge distribution is subject to the Lorentz-FitzGerald contraction, we can turn it into a system with the exact same energy-momentum-mass-velocity relations as the Lorentz-Poincaré electron by adding the appropriate non-electromagnetic stabilizing mechanism.⁵⁰ Of course, as the analysis in this section, based

⁴⁹ Which of the two situations obtains can obviously not be decided experimentally. The rest mass of the electron can be determined but that value can be represented by $m_0 = U_0 / c^2$ or by $m_0 = 4U_0 / 3c^2$ simply by adjusting the radius of the electron, which cannot be determined experimentally.

⁵⁰ This stabilizing system will not be as simple as the Poincaré pressure for the Lorentz-Poincaré electron. Without the spherical symmetry of this specific model, eq. (6.12) for the non-electromagnetic part of the energy-momentum tensor will be more complicated. See (Janssen 1995, sec. 2.3.3, especially eq. (2.96))

on (Laue 1911a), shows, any closed static system will have the same energy-momentum-mass-velocity relations as the Lorentz-Poincaré electron, no matter whether it consists of charges, electromagnetic fields, and Poincaré pressure or of something else altogether. The only thing that matters is that whatever the electron is made of satisfies Lorentz-invariant laws. The restriction to *static* closed systems, moreover, is completely unnecessary. Any closed system will do.⁵¹ In short, there is nothing we can learn about the nature and structure of the electron from studying its energy-momentum-mass-velocity relations.

Lorentz himself emphasized this in lectures he gave at Caltech in 1922. In a section entitled “Structure of the Electron” in the book based on these lectures and published in 1927, he wrote:

The formula for momentum was found by a theory in which it was supposed that in the case of the electron the momentum is determined wholly by that of the electromagnetic field [...] This meant that the whole mass of an electron was supposed to be of electromagnetic nature. Then, when the formula for momentum was verified by experiment, it was thought at first that it was thereby proved that electrons have no “material mass.” Now we can no longer say this. Indeed, the formula for momentum is a general consequence of the principle of relativity, and a verification of that formula is a verification of the principle and tells us nothing about the nature of mass or of the structure of the electron. Therefore physicists are absolutely free to form any hypotheses on the properties and size of electrons that may best suit them. [...] Of course I need hardly mention that, whatever theory we favor, we must suppose that a motion of translation will make the electron contract. Indeed, we want to apply the principle of relativity to the electron also; if then we know what is going on in the electron when it has no motion of translation, we can deduce from the principle in full detail the state that will exist when there is such a motion (Lorentz 1927, 125–126).

for another simple example, the stabilizing mechanism for the surface charge distribution on a plate capacitor, worked out with the help of Tony Duncan.

⁵¹ See the discussion following eq. (2.10) and (Janssen 2003, 46–47).

7. From the electromagnetic view of nature to relativistic continuum mechanics.

Experiment was supposed to be the final arbiter in the debate over the electron models of Abraham, Lorentz-Poincaré, and Bucherer-Langevin. Later analysis, however, showed that none of the experiments of Kaufmann and others were accurate enough to decide between the different models. They only “indicated a large qualitative increase of mass with velocity” (Zahn and Spees 1938, quoted in Miller 1981, 331). All parties involved took these experiments much too seriously, especially when the data favored their own theories. Abraham hyped Kaufmann’s results. Lorentz was too eager to believe Bucherer’s results, while his earlier concern over Kaufmann’s appears to have been somewhat disingenuous. Einstein’s cavalier attitude toward Kaufmann’s experiments stands in marked contrast to his belief in later results purporting to prove him right.

In Abraham’s defense, it should be said that he could also be self-deprecating about his reliance on Kaufmann’s data. At the 78th *Versammlung Deutscher Naturforscher und Ärzte* in Stuttgart in 1906, he got quite a few laughs when he joked: “When you look at the numbers you conclude from them that the deviations from the Lorentz theory are at least twice as big as mine, so you may say that the [rigid] sphere theory represents the reflection of β -rays twice as well as the relativity theory [by which Abraham meant Lorentz’s electron model in this context]” (quoted in Miller 1981, sec. 7.4.3, 221).

In 1906 Lorentz gave a series of lectures at Columbia University in New York that were published in 1909. On the face of it, he seems to have taken Kaufmann’s results quite seriously at the time. He wrote: “His [i.e., Kaufmann’s] new numbers agree within the limits of experimental errors with the formulae given by Abraham, but [...] are decidedly unfavourable to the idea of a contraction such as I attempted to work out” (Lorentz 1915, 212–213; quoted in Miller 1981, sec. 12.4.1). Shortly before his departure for New York, he had told Poincaré the same thing: “Unfortunately my hypothesis of the flattening of electrons is in contradiction with Kaufmann’s results, and I must abandon it. I am therefore at the end of my rope (*au bout de mon latin*).”⁵² These passages strongly suggest that Lorentz took Kaufmann’s results much more seriously than Einstein. Miller indeed draws that conclusion. Lorentz expert A. J. Kox, however, has pointed out to one of us (MJ) that Lorentz’s reaction was probably more ambivalent. This is suggested by what Lorentz continues to say after acknowledging the problem with Kaufmann’s data in

⁵² Lorentz to Poincaré, March 8, 1906 (see Miller 1981, sec. 12.4.1, for the quotation, and pp. 318–319 for a reproduction of the letter in facsimile).

his New York lectures: “Yet, though it seems very likely that we shall have to relinquish this idea altogether, *it is, I think, worth while looking into it somewhat more closely*” (Lorentz 1915, 213; our italics). Lorentz then proceeds to discuss his idea *at length*.

In response to Kaufmann’s alleged refutation of special relativity Einstein wrote in an oft-quoted passage:⁵³ “Abraham’s and Bucherer’s theories of the motion of the electron yield curves that are significantly closer to the observed curve than the curve obtained from the theory of relativity. However, the probability that their theories are correct is rather small, in my opinion, because their basic assumptions concerning ... the moving electron are not suggested by theoretical systems that encompass larger complexes of phenomena” (Einstein 1907b, 439). This is fair enough. When the data went his way, however, Einstein took them much more seriously. In early 1917, Friedrich Adler, detained in Vienna awaiting trial for his assassination of the Austrian prime minister Count Stürgkh in November 1916, began sending Einstein letters and manuscripts attacking special relativity.⁵⁴ He was still at it in the fall of 1918, when the exchange that is interesting for our purposes took place. Einstein wrote: “for a while Bucherer advocated a theory that comes down to a different choice for l [see eq. (3.3) and Fig. 1]. But a different choice for l is out of the question now that the laws of motion of the electron have been verified with great precision.”⁵⁵ From his prison cell in Stein an der Donau Adler replied: “Now, I would be very interested to hear, *which* experiments you see as definitively decisive about the laws of motion of the electron. For as far as my knowledge of the literature goes, I have not found any claim of a final decision.”⁵⁶ Adler

⁵³ See, e.g., (Holton 1988, 252–253), (Miller 1981, sec. 12.4.3), and (Janssen 2002a, 462, note 9).

⁵⁴ See Adler to Einstein, March 9, 1917 (Einstein 1987–2002, Vol. 8, Doc. 307). In 1909 Adler had supported Einstein’s candidacy for a post at the University of Zurich for which both of them had applied (see Einstein to Michele Besso, April 29, 1917 [Einstein 1987–2002, Vol. 8, Doc. 331]). Einstein reciprocated in 1917 by drafting a petition on behalf of a number of Zürich physicists asking the Austrian authorities for leniency in Adler’s case, even as Adler was busying himself with a critique of his benefactor’s theories (see the letter to Besso quoted above). Adler’s father, the well-known Austrian social democrat Victor Adler, considered using his son’s railings against relativity for an insanity defense. His son, however, was determined to stand by his critique of relativity, even if it meant ending up in front of the firing squad. Adler was in fact sentenced to death but it was clear to all involved that he would not be executed. The death sentence was commuted to eighteen years in prison on appeal and Adler was pardoned immediately after the war. This bizarre story is related in (Fölsing 1997, 402–405). For an analysis of the psychology behind Adler’s burning martyrdom, see (Ardelt 1984).

⁵⁵ Einstein to Adler, September 29, 1918 (Einstein 1987–2002, Vol. 8, Doc. 628; translation here and in the following based on Ann M. Hentschel’s).

⁵⁶ Adler to Einstein, October 12, 1918 (Einstein 1987–2002, Vol. 8, Doc. 632; Adler’s emphasis).

went on to quote remarks from Laue, Lorentz, and the experimentalist Erich Hupka, spanning the years 1910–1915, all saying that this was still an open issue.⁵⁷ In his response Einstein cited three recent papers (published between 1914 and 1916), which, he wrote, “have so to speak *conclusively shown* [*sicher bewiesen*] that the relativistic laws of motion of the electron apply (as opposed to, for instance, those of Abraham)” (Einstein’s emphasis).⁵⁸ Even considering the context in which it was made, this is a remarkably strong statement.

Much more interesting than the agreement between theory and experiment or the lack thereof were the theoretical arguments that Abraham and Lorentz put forward in support of their models. Lorentz was right in thinking that it was no coincidence that his contractile electron exhibited exactly the velocity dependence he needed to account for the absence of ether drift (see the discussion following eq. (4.21)). He could not have known at the time that this particular velocity dependence would turn out to be a generic feature of relativistic closed systems. As the quotation at the end of sec. 6 shows, he did recognize this later on. Abraham was right that fast electrons call for a new mechanics. His new electromagnetic mechanics is much closer to relativistic mechanics than to Newtonian mechanics. Like Lorentz, he just did not realize that this new mechanics reflected a new kinematics rather than the electromagnetic nature of all matter. Abraham at least came to accept that Minkowski space-time was the natural setting for his electromagnetic program.

Proceeding along similar lines as Abraham in developing his electromagnetic mechanics, we can easily get from Newtonian particle mechanics to relativistic continuum mechanics and back again. The first step is to read $\mathbf{F} = m\mathbf{a}$ as expressing momentum conservation (cf. the discussion following eq. (2.12) in sec. 2.2). In continuum mechanics, the differential form of the conservation laws is the fundamental law and the integral form is a derived law. In other words, the fundamental conservation laws are expressed in local rather than global terms. This reflects the transition from a particle ontology to a field ontology. Special relativity integrates the laws of momentum and energy conservation. These laws, of course, are Lorentz-invariant rather than Galilean-invariant. We thus arrive at the fundamental law of relativistic continuum mechanics, the Lorentz-invariant differential law of energy-momentum conservation, $\partial_\nu T^{\mu\nu} = 0$. To recap: there are three key elements in the transition from Newtonian

⁵⁷ Cf., however, the quotation from Lorentz in note 64 below.

⁵⁸ Einstein to Adler, October 20, 1918 (Einstein 1987–2002, Vol. 8, Doc. 636).

particle mechanics based on $\mathbf{F} = m\mathbf{a}$ to relativistic continuum mechanics based on $\partial_\nu T^{\mu\nu} = 0$. They are (in no particular order): the transition from Galilean invariance to Lorentz invariance, the focus on conservation laws rather than force laws, and the transition from a particle ontology to a field ontology.

We now show how, once we have relativistic continuum mechanics, we recover Newtonian particle mechanics. Consider a closed system described by continuous (classical) fields such that the total energy-momentum tensor $T_{\text{tot}}^{\mu\nu}$ of the system can be split into a part describing a localizable particle (e.g., an electron a la Lorentz-Poincaré⁵⁹) and a part describing its environment (e.g., an external electromagnetic field):

$$T_{\text{tot}}^{\mu\nu} = T_{\text{particle}}^{\mu\nu} + T_{\text{environment}}^{\mu\nu}. \quad (7.1)$$

Using our fundamental law, $\partial_\nu T_{\text{tot}}^{\mu\nu} = 0$, integrated over space, we find

$$0 = \int \partial_\nu T_{\text{tot}}^{\mu\nu} d^3x = \int \partial_\nu T_{\text{particle}}^{\mu\nu} d^3x + \int \partial_\nu T_{\text{environment}}^{\mu\nu} d^3x. \quad (7.2)$$

As long as $T_{\text{particle}}^{\mu\nu}$ drops off faster than $1/r^2$ as we go to infinity, Gauss' theorem tells us that

$$\int \partial_i T_{\text{particle}}^{i\mu} d^3x = 0. \quad (7.3)$$

For $\partial_\nu T_{\text{environment}}^{\mu\nu}$ we can substitute minus the density f_{external}^μ of the four-force acting on the particle. The spatial components of eq. (7.2) can thus be written as

$$\partial_0 \int T_{\text{particle}}^{i0} d^3x = \int f_{\text{external}}^i d^3x. \quad (7.4)$$

The right-hand side gives the components of $\mathbf{F}_{\text{external}}$. Since $P_{\text{particle}}^\mu \equiv \frac{1}{c} \int T_{\text{particle}}^{\mu 0} d^3x$ and $x^0 = ct$, the left-hand side is the time derivative of the particle's momentum. Eq. (7.4) is thus equivalent to

$$\frac{d\mathbf{P}_{\text{particle}}}{dt} = \mathbf{F}_{\text{external}}. \quad (7.5)$$

⁵⁹ In general we need the fields associated with the particle to be sharply peaked around the worldline of the particle, a four-dimensional 'world-tube.'

This equation has the same form (and the same transformation properties) as Abraham's electromagnetic equation of motion (2.27). In Abraham's equation, $\mathbf{P}_{\text{particle}}$ is the electromagnetic momentum of the electron, and $\mathbf{F}_{\text{external}}$ is the Lorentz force exerted on the electron by the external fields. Under the appropriate circumstances and with the appropriate identification of the Newtonian mass m , Abraham's electromagnetic equation of motion reduces to Newton's second law, $\mathbf{F} = m\mathbf{a}$ (see eq. (2.34)). The same is true for our more general eq. (7.5). This equation, however, is not tied to electrodynamics. It is completely agnostic about the nature of both the particle and the external force. The only thing that matters is that it describes systems in Minkowski space-time, which obey relativistic kinematics. $\mathbf{P}_{\text{particle}}$ and $\mathbf{F}_{\text{external}}$, like Abraham's electromagnetic momentum and the Lorentz force, only transform as vectors under Galilean transformations in the limit of low velocities, where Lorentz transformations are indistinguishable from Galilean transformations. They inherit their transformation properties from $\partial_\nu T_{\text{particle}}^{\mu\nu}$ and f_{external}^μ , respectively, which transform as four-vectors under Lorentz transformations.

It only makes sense to split the total energy-momentum tensor $T_{\text{tot}}^{\mu\nu}$ into a particle part and an environment part, if the interactions holding the particle together are much stronger than the interactions of the particle with its environment. Typically, therefore, the energy-momentum of the particle taken by itself will very nearly be conserved, i.e.,

$$\partial_\nu T_{\text{particle}}^{\mu\nu} \approx 0. \quad (7.6)$$

This means that the particle's four-momentum will to all intents and purposes transform as a four-vector under Lorentz transformations and satisfy the relations for a strictly closed system (see eqs. (2.1)–(2.10)):

$$P_{\text{particle}}^\mu \equiv \frac{1}{c} \int T_{\text{particle}}^{\mu 0} d^3x \approx (\gamma m_0 c, \gamma m_0 \mathbf{v}). \quad (7.7)$$

Inserting $\mathbf{P}_{\text{particle}} = \gamma m_0 \mathbf{v}$ into eq. (7.5), we can reduce the problem in relativistic continuum mechanics that we started from in eq. (7.1) to a problem in the relativistic mechanics of point particles. In the limit of small velocities, such problems once again reduce to problems in the Newtonian mechanics of point particles.

To the best of our knowledge, this way of recovering particle mechanics from what might be called 'field mechanics' was first worked out explicitly in the context of general

rather than special relativity (Einstein 1918, Klein 1918).⁶⁰ Relativistic continuum mechanics played a crucial role in the development of general relativity. For one thing, the energy-momentum tensor is the source of the gravitational field in general relativity.⁶¹ Even before the development of general relativity, Einstein recognized the importance of relativistic continuum mechanics. In an unpublished manuscript of 1912, he wrote:

The general validity of the conservation laws and of the law of the inertia of energy [...] suggest that [the symmetric energy-momentum tensor $T^{\mu\nu}$ and the equation $f^\mu = -\partial_\nu T^{\mu\nu}$] are to be ascribed a general significance, even though they were obtained in a very special case [i.e., electrodynamics]. We owe this generalization, *which is the most important new advance in the theory of relativity*, to the investigations of Minkowski, Abraham, Planck, and Laue (Einstein 1987–2002, Vol. 4, Doc. 1, [p. 63]; our emphasis).

Einstein went on to give a clear characterization of relativistic continuum mechanics:

To every kind of material process we want to study, we have to assign a symmetric tensor ($T_{\mu\nu}$) [...] Then [$f^\mu = -\partial_\nu T^{\mu\nu}$] must always be satisfied. The problem to be solved always consists in finding out how ($T_{\mu\nu}$) is to be formed from the variables characterizing the processes under consideration. If several processes can be isolated in the energy-momentum balance that take place in the same region, we have to assign to each individual process its own stress-energy tensor ($T_{\mu\nu}^{(1)}$), etc., and set ($T_{\mu\nu}$) equal to the sum of these individual tensors (ibid.).

As the development of general theory of relativity was demonstrating the importance of continuum mechanics, developments in quantum theory—the Bohr model and Sommerfeld’s relativistic corrections to it—rehabilitated particle mechanics, be it of the Newtonian or of the relativistic variety. As a result, relativistic continuum mechanics

⁶⁰ Einstein and Klein corresponded about this issue in 1918 (Einstein 1987–2002, Vol. 8, Docs. 554, 556, 561, 566, and 581). See also Hermann Weyl to Einstein, 16 November 1918 (Einstein 1987–2002, Vol. 8, Doc. 657). A precursor to this approach can be found in (Einstein and Grossmann 1913, sec. 4), where Einstein pointed out that the geodesic equation, which governs the motion of a test particle in a gravitational field, can be obtained by integrating $T^{\mu\nu}_{; \nu} = 0$ —the vanishing of the covariant divergence of $T^{\mu\nu}$, the generalization of $\partial_\nu T^{\mu\nu} = 0$ in general relativity—over the ‘worldtube’ of the corresponding energy-momentum tensor for pressureless dust (“thread of flow” [*Stromfaden*] is the term Einstein used). This argument can also be found in the so-called Zurich Notebook (Einstein 1987–2002, Vol. 4, Doc. 10, [p. 10] and [p. 58]). For analysis of these passages, see “A Commentary on Einstein’s Zurich Notebook” in (Renn et al. forthcoming, sec. 3 and 5.5.10; the relevant pages of the notebook are referred to as ‘5R’ and ‘43L’).

⁶¹ See (Renn and Sauer forthcoming) for extensive discussion of the role of the energy-momentum tensor in the research that led to general relativity.

proved less important for subsequent developments in areas of physics other than general relativity than Einstein thought in 1912 and than our analysis in this paper suggests. The key factor in this was that it gradually became clear in the 1920s that elementary particles are point-like and not spatially extended like the electron models discussed in this paper. That special relativity precludes the existence of rigid bodies is just one of the problems such models are facing.

In hindsight, Lorentz, the guarded Dutchman, comes out looking much better than Abraham, his impetuous German counterpart. At one point, for instance, Lorentz (1915, 215) cautioned:

In speculating on the structure of these minute particles we must not forget that there may be many possibilities not dreamt of at present; it may very well be that other internal forces serve to ensure the stability of the system, and perhaps, after all, we are wholly on the wrong track when we apply to the parts of an electron our ordinary notion of force (Lorentz 1915, 215; quoted approvingly in Pais 1972, 83).

Even a crude operationalist argument of the young Wolfgang Pauli, which would have made his godfather Ernst Mach proud, can look prescient in hindsight. Criticizing the work of later proponents of the electromagnetic worldview in his review article on relativity, Pauli concluded:

Finally, a conceptual doubt should be mentioned. The continuum theories make direct use of the ordinary concept of electric field strength, even for the fields in the interior of the electron. This field strength, however, is defined as the force acting on a test particle, and since there are no test particles smaller than an electron or a hydrogen nucleus the field strength at a given point in the interior of such a particle would seem to be unobservable by definition, and thus be fictitious and without physical meaning (Pauli 1921, 206).

This moved Valentin Bargmann (1960, 189)—who had accompanied Einstein on his quest for a classical unified field theory, a quest very much in the spirit of Abraham’s electromagnetic program—to write in the Pauli memorial volume:

A physicist will feel both pride and humility when he reads Pauli’s remarks today. In the light of our present knowledge the attempts which Pauli criticizes may seem hopelessly naïve, although it was certainly sound practice to investigate what the profound new ideas of general relativity would contribute to the understanding of the thorny problem of matter (Bargmann 1960, 189).

We conclude our paper by quoting and commenting on two oft-quoted passages that nicely illustrate some of the key points of our paper. The first is a brief exchange between

Planck and Abraham⁶² following a lecture by the former at the *Naturforscherversammlung* in Stuttgart on September 19, 1906. Planck talked about “[t]he Kaufmann measurements of the deflectability of β -rays and their relevance for the dynamics of electrons.” Abraham, Bucherer,⁶³ Kaufmann, and Sommerfeld all took part in the discussion afterwards. It was Planck who got to the heart of the matter:

Abraham is right when he says that the essential advantage of the sphere theory would be that it be a purely electrical theory. If this were feasible, it would be very beautiful indeed, but for the time being it is just a postulate. At the basis of the Lorentz-Einstein theory lies another postulate, namely that no absolute translation can be detected. These two postulates, it seems to me, cannot be combined, and what it comes down to is which postulate one prefers. My sympathies actually lie with the Lorentzian postulate (Planck 1906b, 761).

In response Sommerfeld, pushing forty at the time, quipped: “I suspect that the gentlemen under forty will prefer the electrodynamical postulate, while those over forty will prefer the mechanical-relativistic postulate” (Ibid.). The reaction of the assembled physicists to Sommerfeld’s quick retort has also been preserved in the transcript of this session: “laughter” (*Heiterkeit*). This exchange between Planck and Sommerfeld is perhaps the clearest statement in the contemporary literature of the dilemma that lies behind the choice between the electron models of Abraham and Lorentz. Physicists had to decide what they thought was more important, full relativity of uniform motion or the reduction of mechanics to electrodynamics. We find it very telling that in 1906 a leader in the field such as Sommerfeld considered special relativity to be the conservative and the electromagnetic program to be the progressive option! Only a few years later, Sommerfeld jumped ship and joined the relativity camp.

The second passage that we want to look at comes from Lorentz’s important book *The Theory of Electrons*, based on his 1906 lectures in New York and first published in 1909. Referring to Einstein and special relativity, Lorentz wrote

His results concerning electromagnetic and optical phenomena (leading to the same contradiction with Kaufmann’s results that was pointed out in §179^[64]) agree in the main

⁶² This exchange is also discussed, for instance, in (Miller 1981, sec. 7.3.4), (McCormmach 1970, 489–490), and (Jungnickel and McCormmach 1986, 249–250).

⁶³ Understandably, Bucherer took exception to the fact that Planck only discussed the electron models of Lorentz and Abraham (Planck 1906, 760).

⁶⁴ In the second edition, Lorentz added the following footnote at this point: “Later experiments [...] have confirmed [eq. 2.43] for the transverse electromagnetic mass, so that, in all probability, the only objection

with those which we have obtained in the preceding pages, the chief difference being that Einstein simply postulates what we have deduced, with some difficulty and not altogether satisfactorily, from the fundamental equations of the electromagnetic field. (Lorentz 1915, 229–230).

The parenthetical reference to “Kaufmann’s results” suggests that the famous clause that concludes this sentence—“Einstein simply postulates what we have deduced [...] from the fundamental equations of the electromagnetic field”—refers, at least in part, to Lorentz’s own struggles with the velocity dependence of electron mass.⁶⁵ The relativistic derivation of these relations is mathematically equivalent to Lorentz’s 1899 derivation, which showed that they are necessary to render ether drift unobservable (see sec. 3, eqs. (3.8)–(3.14)). From Lorentz’s point of view, the relativistic derivation therefore amounted to nothing more than postulating these relations on the basis of the relativity principle. Lorentz himself had gone beyond showing that these relations follow from the requirement, formally identical to the relativity principle, that ether drift can never be detected. He had gone to the trouble of producing a concrete model of the electron such that its mass exhibited exactly the desired velocity-dependence (see sec. 4, eqs. (4.16)–(4.22)). As we saw at the end of sec. 6, by 1922, if not much earlier, Lorentz had recognized that this had led him on a wild goose chase: “the formula for momentum [from which those for the velocity dependence of mass are a direct consequence] is a general consequence of the principle of relativity, and a verification of that formula is a verification of the principle and tells us nothing about the nature of mass or of the structure of the electron.” This was Lorentz’s way of saying what Pais said in the quotation with which we began this paper.

that could be raised against the hypothesis of the deformable electron and the principle of relativity has now been removed” (Lorentz 1915, 339).

⁶⁵ For more extensive discussion of this passage, see (Janssen 1995, secs. 4.3).

Acknowledgments: We want to thank Tony Duncan, Gordon Fleming, A. J. Kox, Christoph Lehner, John Norton, Jürgen Renn, Serge Rudaz, Robert Schulmann, and John Stachel for helpful discussions, suggestions, and references. One of us (MJ) gratefully acknowledges support from the *Max-Planck-Institut für Wissenschaftsgeschichte* in Berlin.

References

- Abraham, Max. 1902a. “Dynamik des Elektrons.” *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 20–41.
- — —. 1902b. “Prinzipien der Dynamik des Elektrons.” *Physikalische Zeitschrift* 4: 57–63.
- — —. 1903. “Prinzipien der Dynamik des Elektrons.” *Annalen der Physik* 10: 105–179.
- — —. 1904. “Die Grundhypothesen der Elektronentheorie.” *Physikalische Zeitschrift* 5: 576–579.
- — —. 1905. *Theorie der Elektrizität*. Vol. 2. *Elektromagnetische Theorie der Strahlung*. Leipzig: Teubner.
- — —. 1909. “Zur Elektrodynamik bewegter Körper.” *Circolo Matematico di Palermo. Rendiconti* 28: 1–28.
- Ardelt, Rudolf G. 1984. *Friedrich Adler. Probleme einer Persönlichkeitsentwicklung um die Jahrhundertwende*. Vienna: Österreichischer Bundesverlag.
- Balazs, N. L. 1972. “The Acceptability of Physical Theories: Poincaré versus Einstein.” Pp. 21–34 in *General Relativity. Papers in Honour of J. L. Synge*. Edited by L. O’Raifeartaigh. Oxford: Clarendon Press.
- Bargmann, Valentin. 1960. “Relativity.” Pp. 187–198 in *Theoretical Physics in the Twentieth Century. A Memorial Volume to Wolfgang Pauli*. Edited by M. Fierz and V. F. Weisskopf. New York: Interscience Publishers.
- Bucherer, Alfred H. 1904. *Mathematische Einführung in die Elektronentheorie*. Leipzig: Teubner.
- — —. 1905. “Das deformierte Elektron und die Theorie des Elektromagnetismus.” *Physikalische Zeitschrift* 6: 833–834.
- — —. 1907. “On a New Principle of Relativity in Electromagnetism.” *Philosophical Magazine* 13: 413–420.

- — —. 1908a. “On the Principle of Relativity and on the Electromagnetic Mass of the Electron. A Reply to Mr. E. Cunningham.” *Philosophical Magazine* 15: 316–318. Response to (Cunningham 1907).
- — —. 1908b. “On the Principle of Relativity. A Reply to Mr. E. Cunningham.” *Philosophical Magazine* 16: 939–940. Response to (Cunningham 1908).
- — —. 1908c. “Messungen an Becquerelstrahlen. Die experimentelle Bestätigung der Lorentz-Einsteinschen Theorie.” *Physikalische Zeitschrift* 9: 755–762.
- Butler, J. W. 1968. “On the Trouton-Noble experiment.” *American Journal of Physics* 36: 936–941.
- Corry, Leo. 1997 “Hermann Minkowski and the Postulate of Relativity.” *Archive for History of Exact Sciences* 51: 273–314.
- — —. 1999. “From Mie’s Electromagnetic Theory of Matter to Hilbert’s Unified Foundations of Physics.” *Studies in History and Philosophy of Modern Physics* 30: 159–183.
- Cattani, Carlo, and Michelangelo De Maria. 1989. “The 1915 Epistolary Controversy Between Einstein and Tullio Levi-Civita.” Pp. 175–200 in *Einstein and the History of General Relativity*. Edited by Don Howard and John Stachel. Boston: Birkhäuser.
- Cunningham, Ebenezer. 1907. “On the Electromagnetic Mass of a Moving Electron.” *Philosophical Magazine* 14: 538–547.
- — —. 1908. “On the Principle of Relativity and the Electromagnetic Mass of the Electron. A Reply to Dr. A. H. Bucherer.” *Philosophical Magazine* 16: 423–428. Response to (Bucherer 1908a).
- Cushing, James T. 1981. “Electromagnetic Mass, Relativity, and the Kaufmann Experiments.” *American Journal of Physics* 49: 1133–1149.
- Damerow, Peter, Gideon Freudenthal, Peter McLaughlin, and Jürgen Renn. 2004. *Exploring the Limits of Preclassical Mechanics. A Study of Conceptual Development in Early Modern Science: Free Fall and Compounded Motion in the Work of Descartes, Galileo, and Beeckman*. 2nd ed. New York: Springer.
- Darrigol, Olivier. 1995. “Henri Poincaré’s Criticism of Fin de Siècle Electrodynamics.” *Studies in History and Philosophy of Modern Physics* 26:1–44.
- — —. 2000. *Electrodynamics from Ampère to Einstein*. Oxford: Oxford University Press.
- Earman, John. 2003. “The Cosmological Constant, the Fate of the Universe, Unimodular Gravity, and all that.” *Studies in History and Philosophy of Modern Physics* 34: 559–577.

- Ehrenfest, Paul. 1906. “Zur Stabilitätsfrage bei den Bucherer-Langevin-Elektronen.” *Physikalische Zeitschrift* 7: 302–303
- — —. 1907. “Die Translation deformierbarer Elektronen und der Flächensatz.” *Annalen der Physik* 23: 204–205.
- Einstein, Albert. 1905. “Zur Elektrodynamik bewegter Körper.” *Annalen der Physik* 17: 891–921. Reprinted in (Einstein 1987–2002), Vol. 2, Doc. 23.
- — —. 1907a. “Bemerkungen zu der Notiz von Hrn. Paul Ehrenfest: „Die Translation deformierbarer Elektronen und der Flächensatz.“” *Annalen der Physik* 23: 206–208. Reprinted in (Einstein 1987–2002), Vol. 2, Doc. 44.
- — —. 1907b. “Über die vom Relativitätsprinzip geforderte Trägheit der Energie.” *Annalen der Physik* 23: 371–384. Reprinted in (Einstein 1987–2002), Vol. 2, Doc. 45.
- — —. 1918. “Der Energiesatz in der allgemeinen Relativitätstheorie.” *Preußische Akademie der Wissenschaften. Sitzungsberichte* 448–459. Reprinted in (Einstein 1987–2002), Vol. 7, Doc. 9.
- — —. 1919. “Spielen Gravitationsfelder im Aufbau der materiellen Elementarteilchen eine wesentliche Rolle?” *Königlich Preußische Akademie der Wissenschaften (Berlin). Sitzungsberichte*: 349–356. Reprinted in (Einstein 1987–2002), Vol. 7, Doc. 17.
- — —. 1987–2002. *The Collected Papers of Albert Einstein*. 8 Vols. Edited by John Stachel et al. Princeton: Princeton University Press.
- Einstein, Albert, and Marcel Grossmann. 1913. *Entwurf einer verallgemeinerten Relativitätstheorie und einer Theorie der Gravitation*. Leipzig: Teubner. Reprinted in (Einstein 1987–2002), Vol. 4, Doc. 13.
- Fermi, Enrico. 1921. “Sulla dinamica di un sistema rigido di cariche elettriche in moto traslatorio” *Nuovo Cimento* 22: 199–207. Reprinted in (Fermi 1962–1965, 1–7).
- — —. 1922. “Über einen Widerspruch zwischen der elektrodynamischen und der relativistischen Theorie der elektromagnetischen Masse.” *Physikalische Zeitschrift* 23: 340–344.
- — —. 1962–1965. *The Collected Works of Enrico Fermi*. 2 Vols. Edited by Emilio Segrè. Chicago, Rome: University of Chicago Press, Accademia Nazionale dei Lincei.
- Feynman, Richard P., Robert B. Leighton, and Matthew Sands. 1964. *The Feynman Lectures on Physics*. Vol. II. *Mainly Electromagnetism and Matter*. Reading, MA: Addison-Wesley.
- Galison, Peter L. 1979. “Minkowski’s Space-Time: From Visual Thinking to the Absolute World.” *Historical Studies in the Physical Sciences* 10: 85–121.

- Goldberg, Stanley. 1970. "The Abraham Theory of the Electron: The Symbiosis of Experiment and Theory." *Archive for History of Exact Sciences* 7: 7–25.
- Griffith, David J. 1999. *Introduction to Electrodynamics*. 3rd ed. Upper Saddle River, New Jersey: Prentice Hall.
- Grøn, Ø. 1985. "Repulsive Gravitation and Electron Models." *Physical Review D* 31: 2129–2131.
- — —. 1988. "Poincaré Stress and the Reissner-Nordström Repulsion." *General Relativity and Gravitation* 20: 123–129.
- Herglotz, Gustav. 1903. "Zur Elektronentheorie" *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 357–382.
- — —. 1910. "Über den vom Standpunkt des Relativitätsprinzips aus als 'starr' zu bezeichnenden Körper." *Annalen der Physik* 31: 393–415.
- — —. 1911. "Über die Mechanik des deformierbaren Körpers vom Standpunkte der Relativitätstheorie." *Annalen der Physik* 36: 493–533.
- Holton, Gerald. 1988. "Mach, Einstein, and the Search for Reality." Pp. 237–277 in *Thematic Origins of Scientific Thought: Kepler to Einstein*. Rev. ed. Cambridge: Harvard University Press.
- Hon, Giora. 1995. "Is the Identification of Experimental Error Contextually Dependent? The Case of Kaufmann's Experiment and Its Varied Reception." Pp. 170–223 in *Scientific Practice. Theories and Stories of Doing Physics*. Edited by Jed Z. Buchwald. Chicago: University of Chicago Press.
- Jackson, John D. 1975. *Classical Electrodynamics*. 2nd ed. New York: John Wiley & Sons.
- Jammer, Max. 1997. *Concepts of Mass in Classical and Modern Physics*. New York: Dover.
- Janssen, Michel. *A Comparison between Lorentz's Ether Theory and Einstein's Special Theory of Relativity in the Light of the Experiments of Trouton and Noble*. Ph.D. Thesis. University of Pittsburgh, 1995.
- — —. 2002a. "COI Stories: Explanation and Evidence in the History of Science." *Perspectives on Science* 10: 457–522.
- — —. 2002b. "Reconsidering a Scientific Revolution: the Case of Lorentz versus Einstein." *Physics in Perspective* 4: 421–446.
- — —. 2003. "The Trouton Experiment, $E = mc^2$, and a Slice of Minkowski Space-Time." Pp. 27–54 in *Revisiting the Foundations of Relativistic Physics: Festschrift in Honor of John Stachel*. Edited by Abhay Ashtekar et al. (ed.). Dordrecht: Kluwer.

- Janssen, Michel, and John Stachel. 2004. *The Optics and Electrodynamics of Moving Bodies*. Preprint 265. Max Planck Institute for History of Science, Berlin.
- Jungnickel, Christa, and Russell McCormach. 1986. *Intellectual Mastery of Nature: Theoretical Physics from Ohm to Einstein*. Vol. 2. *The Now Mighty Theoretical Physics, 1870-1925*. Chicago: University of Chicago Press.
- Kaufmann, Walter. 1901a. "Die Entwicklung des Elektronenbegriffs." Pp. 115–126 in *Verhandlungen der Gesellschaft Deutscher Naturforscher und Ärzte*. 73. *Versammlung zu Hamburg*. 22.–28. September 1901. Part 1. Edited by Albert Wangering. Leipzig: Vogel. Reprinted in *Physikalische Zeitschrift* 3: 9–15.
- — —. 1901b. "Methode zur exakten Bestimmung von Ladung und Geschwindigkeit der Becquerelstrahlen." *Physikalische Zeitschrift* 2: 602–603.
- — —. 1901c. "Die magnetische und elektrische Ablenkbarkeit der Becquerelstrahlen und die scheinbare Masse der Elektronen." *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 143–155.
- — —. 1902a. "Über die elektromagnetische Masse des Elektrons." *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 291–303.
- — —. 1902b. "Die elektromagnetische Masse des Elektrons." *Physikalische Zeitschrift* 4: 54–57.
- — —. 1903. "Über die 'elektromagnetische Masse' der Elektronen." *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 90–103.
- — —. 1905. "Über die Konstitution des Elektrons." *Königlich Preußische Akademie der Wissenschaften (Berlin). Sitzungsberichte* 949–956.
- — —. 1906a. "Über die Konstitution des Elektrons." *Annalen der Physik* 19: 487–553.
- — —. 1906b. "Nachtrag zu der Abhandlung „Über die Konstitution des Elektrons“." *Annalen der Physik* 20: 639–640.
- — —. 1907. "Bemerkungen zur Herrn Plancks „Nachtrag zu der Besprechung der Kaufmannschen Ablenkungsmessungen“." *Deutsche Physikalische Gesellschaft. Verhandlungen* 9: 667–673.
- Keswani, G. H., and C. W. Kilmister. 1983. "Intimations of Relativity: Relativity Before Einstein." *British Journal for the Philosophy of Science* 34: 343–354.
- Kilmister, C. W., ed. 1970. *Special Theory of Relativity*. London: Pergamon.

- Klein, Felix. 1918. "Über die Integralform der Erhaltungssätze und die Theorie der räumlich-geschlossenen Welt." *Königliche Gesellschaft der Wissenschaften zu Göttingen. Nachrichten* 394–423.
- Kragh, Helge. 1999. *Quantum Generations. A History of Physics in the Twentieth Century*. Princeton: Princeton University Press.
- Langevin, Paul. 1905. "La physique des électrons." *Revue générale des sciences pures et appliquées* 16: 257–276.
- Laue, Max. 1911a. "Zur Dynamik der Relativitätstheorie." *Annalen der Physik* 35 (1911): 524–542. Reprinted in (Laue 1961), Vol. 1, pp. 135–153.
- — —. 1911b. *Das Relativitätsprinzip*. Braunschweig: Vieweg.
- Laue, Max von. 1961. *Gesammelte Schriften und Vorträge*. 3 Vols. Braunschweig: Friedrich Vieweg und Sohn.
- Levi-Civita, Tullio. 1907. "Sulla massa elettromagnetica." *Nuovo Cimento* 16: 387–412. Reprinted in (Levi-Civita 1956), Vol. 2, pp. 587–613.
- — —. 1909. "Teoria asintotica delle radiazioni elettriche." *Rendiconti della Reale Accademia dei Lincei* 18: 83–93. Reprinted in (Levi-Civita 1956), Vol. 3, pp. 81–92.
- — —. 1956. *Opere Matematiche. Memorie e Note*. Edited by Francesco Giordani et al. 4 Vols. Bologna: Nicola Zanichelli Editore.
- Lorentz, Hendrik Antoon. 1895. *Versuch einer Theorie der elektrischen und optischen Erscheinungen in bewegten Körpern*. Leiden: Brill. Reprinted in (Lorentz 1934–39), Vol. 5, 1–138.
- — —. 1899. "Vereenvoudigde theorie der elektrische en optische verschijnselen in lichamen die zich bewegen." *Koninklijke Akademie van Wetenschappen te Amsterdam. Wis- en Natuurkundige Afdeeling. Verslagen van de Gewone Vergaderingen* 7 (1898–99): 507–522. Slightly revised English translation: "Simplified theory of electrical and optical phenomena in moving bodies." *Koninklijke Akademie van Wetenschappen te Amsterdam. Section of Sciences. Proceedings* 1 (1898–99): 427–442. Slightly revised French translation: "Théorie simplifiée des phénomènes électriques et optiques dans des corps en mouvement." *Archives Néerlandaises des Sciences Exactes et Naturelles* 7 (1902): 64–80. French translation reprinted in (Lorentz 1934–39), Vol. 5, 139–155.
- — —. 1900. "Über die scheinbare Masse der Elektronen." *Physikalische Zeitschrift* 2 (1900–1901): 78–80 (79–80: discussion). Reprinted in (Lorentz 1934–39), Vol. 3, 113–116.
- — —. 1904a. "Weiterbildung der Maxwellschen Theorie. Elektronentheorie." Pp. 145–288 in *Encyclopädie der mathematischen Wissenschaften, mit Einschluss ihrer Anwendungen*, Vol. 5, *Physik*, Part 2. Edited by Arnold Sommerfeld. Leipzig: Teubner, 1904–1922.

- — — . 1904b. “Electromagnetische verschijnselen in een stelsel dat zich met willekeurige snelheid, kleiner dan die van het licht, beweegt.” *Koninklijke Akademie van Wetenschappen te Amsterdam. Wis- en Natuurkundige Afdeeling. Verslagen van de Gewone Vergaderingen* 12 (1903–04): 986–1009. Translation: “Electromagnetic Phenomena in a System Moving with Any Velocity Smaller Than That of Light.” *Koninklijke Akademie van Wetenschappen te Amsterdam. Section of Sciences. Proceedings* 6 (1903–04): 809–831. Translation reprinted in (Lorentz 1934–39), Vol. 5, 172–197.
- — — . 1905. *Ergebnisse und Probleme der Elektronentheorie. Vortrag, gehalten am 20. Dezember 1904 im Elektrotechnischen Verein zu Berlin*. Berlin: Springer. Reprinted in (Lorentz 1934–39), Vol. 8, 76–124.
- — — . 1915. *The Theory of Electrons and Its Applications to the Phenomena of Light and Radiant Heat*. 2d ed. Leipzig: Teubner, 1915.
- — — . 1927. *Problems of Modern Physics. A Course of Lectures Delivered in the California Institute of Technology*. Boston: Ginn.
- — — . 1934–39. *Collected Papers*. Edited by Pieter Zeeman and Adriaan D. Fokker. 9 Vols. The Hague: Nijhoff.
- McCormmach, Russell. 1970. “H. A. Lorentz and the Electromagnetic View of Nature.” *Isis* 61: 459–497.
- Mie, Gustav. 1912a. “Grundlagen einer Theorie der Materie. (Erste Mitteilung.)” *Annalen der Physik* 37: 511–534.
- — — . 1912b. “Grundlagen einer Theorie der Materie. (Zweite Mitteilung.)” *Annalen der Physik* 39: 1–40.
- — — . 1913. “Grundlagen einer Theorie der Materie. (Dritte Mitteilung, Schluß.)” *Annalen der Physik* 40: 1–66.
- Miller, Arthur I. 1973. “A study of Henri Poincaré’s “Sur la dynamique de l’électron.”” *Archive for the History of Exact Sciences* 10 (1973): 207–328. Reprinted in (Miller 1986).
- — — . 1981. *Albert Einstein’s Special Theory of Relativity. Emergence (1905) and Early Interpretation (1905–1911)*. Reading, Mass.: Addison-Wesley, 1981. Reprinted in 1998 (New York: Springer). Page references are to this reprint.
- — — . 1986. *Frontiers of Physics: 1900–1911*. Boston, Birkhäuser.
- Minkowski, Hermann. 1908. “Die Grundgleichungen für die elektromagnetischen Vorgänge in bewegten Körpern.” *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 53–111. Reprinted in (Minkowski 1967), Vol. 2, 352–404.
- — — . 1909. “Raum und Zeit.” *Physikalische Zeitschrift* 20: 104–111.

- — —. 1967. *Gesammelte Abhandlungen von Hermann Minkowski*. Edited by David Hilbert. New York: Chelsea. Originally published in two volumes, Leipzig, 1911.
- Norton, John D. 1992. “Einstein, Nordström and the Early Demise of Scalar, Lorentz-Covariant Theories of Gravitation.” *Archive for the History of Exact Sciences* 45: 17–94.
- Pais, Abraham, 1972. “The Early History of the Theory of the Electron: 1897–1947.” Pp. 79–93 in *Aspects of Quantum Theory*. Edited by Abdus Salam and Eugene P. Wigner. Cambridge: Cambridge University Press.
- — —. 1982. ‘*Subtle is the Lord ...*’ *The Science and the Life of Albert Einstein*. Oxford: Oxford University Press.
- Pauli, Wolfgang. 1921. “Relativitätstheorie.” Pp. 539–775 in *Encyklopädie der mathematischen Wissenschaften, mit Einschluß ihrer Anwendungen*. Vol. 5, *Physik*, Part 2. Edited by Arnold Sommerfeld. Leipzig: Teubner, 1904–1922. Translation with introduction and supplementary notes: *Theory of Relativity*. G. Field, trans. London: Pergamon, 1958. Reprinted: New York: Dover, 1981. German original (plus German translations of introduction and supplementary notes of English edition) reprinted as *Relativitätstheorie*. Edited by Domenico Giulini. Berlin: Springer, 2000. Page references are to English translation.
- Planck, Max. 1906a. “Das Prinzip der Relativität und die Grundgleichungen der Mechanik.” *Deutsche Physikalische Gesellschaft. Verhandlungen* 8: 136–141. Reprinted in (Planck 1958), Vol. 2, pp. 115–120.
- — —. 1906b. “Die Kaufmannschen Messungen der Ablenkbarkeit der β -Strahlen in ihrer Bedeutung für die Dynamik der Elektronen.” *Deutsche Physikalische Gesellschaft. Verhandlungen* 8: 418–432. Reprinted in *Physikalische Zeitschrift* 7 (1906): 753–759 (760–761: discussion); and in (Planck 1958), Vol. 2, pp. 121–135.
- — —. 1908. “Bemerkungen zum Prinzip der Aktion und Reaktion in der allgemeinen Dynamik.” *Deutsche Physikalische Gesellschaft. Verhandlungen* 6: 728–732. Reprinted in (Planck 1958)
- — —. 1958. *Physikalische Abhandlungen und Vorträge*. 3 Vols. Braunschweig: Friedrich Vieweg und Sohn.
- Poincaré, Henri. 1900. “La théorie de Lorentz et le principe de réaction.” *Archives Néerlandaises des Sciences Exactes et Naturelles* 2:252–278. Reprinted in (Poincaré 1934–54), Vol. 9, 464–493.
- — —. 1905. “Sur la dynamique de l’électron.” *Comptes Rendus de l’Académie des Sciences* 140: 1504–1508. Short version of Poincaré 1906. Reprinted in (Poincaré 1934–54), Vol. 9, 489–493. English translation in (Keswani and Kilmister 1983).
- — —. 1906. “Sur la dynamique de l’électron.” *Rendiconti del Circolo Matematico di Palermo* 21: 129–175. Reprinted in (Poincaré 1934–54), Vol. 9, 494–550. Page

- references are to this reprint. English translation of secs. 1–4 and 9 in (Kilmister 1970), and of secs. 6–8 in (Miller 1973).
- — — . 1934–54. *Œuvres de Henri Poincaré*. 11 Vols. Paris: Gauthiers-Villars.
- Pyenson, Lewis. 1985. *The Young Einstein. The Advent of Relativity*. Bristol and Boston: Adam Hilger.
- Renn, Jürgen, Michel Janssen, Tilman Sauer, John Norton, and John Stachel. Forthcoming. *The Genesis of General Relativity: Documents and Interpretation*. Vol. 1. *General Relativity in the Making: Einstein’s Zurich Notebook*. Dordrecht: Kluwer.
- Renn, Jürgen, and Tilman Sauer. Forthcoming. “A Pathway out of Classical Physics.” To appear in (Renn et al. forthcoming).
- Rohrlich, Fritz. 1960. “Self-energy and the Stability of the Classical Electron.” *American Journal of Physics* 28: 639–643.
- — — . 1965. *Classical Charged Particles: Foundations of Their Theory*. Reading, MA: Addison-Wesley.
- — — . 1973. “The Electron: Development of the First Elementary Particle Theory.” . Pp. 331–369 in *The Physicist’s Conception of Nature*. Edited by Jagdish Mehra. Dordrecht: Reidel.
- Schwartz, H. M. 1971. “Poincaré’s Rendiconti Paper on Relativity. Part I.” *American Journal of Physics* 39: 1287–1294.
- — — . 1972. “Poincaré’s Rendiconti Paper on Relativity. Part II.” *American Journal of Physics* 40: 862–872.
- Schwarzschild, Karl. 1903a. “Zur Elektrodynamik. I. Zwei Formen des Prinzips der kleinsten Action in der Elektronentheorie.” *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 126–131.
- — — . 1903b. “Zur Elektrodynamik. II. Die elementare elektrodynamische Kraft.” *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 132–141.
- — — . 1903c. “Zur Elektrodynamik. III. Ueber die Bewegung des Elektrons.” *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 245–278.
- Segrè, Emilio. 1970. *Enrico Fermi. Physicist*. Chicago: University of Chicago Press.
- Sommerfeld, Arnold. 1904a. “Zur Elektronentheorie. I. Allgemeine Untersuchung des Feldes eines beliebig bewegten Elektrons.” *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 99–130.

- — — . 1904b. “Zur Elektronentheorie. II. Grundlage für eine allgemeine Dynamik des Elektrons.” *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 363–439.
- — — . 1905. “Zur Elektronentheorie. III. Ueber Lichtgeschwindigkeits- und Ueberlichtgeschwindigkeits-Elektronen.” *Königliche Gesellschaft der Wissenschaften zu Göttingen. Mathematisch-physikalische Klasse. Nachrichten*: 201–235.
- — — . 1910a. “Zur Relativitätstheorie I. Vierdimensionale Vektoralgebra.” *Annalen der Physik* 32: 749–776.
- — — . 1910b. “Zur Relativitätstheorie II. Vierdimensionale Vektoranalysis.” *Annalen der Physik* 33: 649–689.
- Teukolsky, Saul A. 1996. “The explanation of the Trouton-Noble experiment revisited.” *American Journal of Physics* 64: 1104–1106.
- Trouton, Frederick T., and Henry R. Noble. 1903. “The Mechanical Forces Acting On a Charged Electric Condenser Moving Through Space.” *Philosophical Transactions of the Royal Society* 202: 165–181.
- Warwick, Andrew. 2003. *Masters of Theory. Cambridge and the Rise of Mathematical Physics*. Chicago: University of Chicago Press.
- Wien, Wilhelm. 1900. “Über die Möglichkeit einer elektromagnetischen Begründung der Mechanik.” *Archives Néerlandais des sciences exactes et naturelles* 2: 96–107. Reprinted in *Annalen der Physik* 5 (1901): 501–513.
- Yaghjian, Arthur D. 1992. *Relativistic Dynamics of a Charged Sphere. Updating the Lorentz-Abraham Model*. Berlin: Springer.
- Zahar, Elie. 1989. *Einstein’s Revolution: a Study in Heuristic*. La Salle, IL: Open Court.
- Zahn, C.T. and A. A. Spees. 1938. “A Critical Analysis of the Classical Experiments On the Variation of Electron Mass.” *Physical Review* 53: 511–521.